1	DRAFT AOAC SMPR 2016.XXX; Version 4; December 9, 2015.		
2 3 4 5	Sta	indard Method Performance Requirements for Determination of Ethanol in Kombucha	
5 6 7	Int	ended Use: Use by trained technicians in a laboratory for routine quality assurance testing.	
8 9 10 11	1.	Applicability : Determination of low levels of ethanol as expressed as alcohol by volume (ABV) in kombucha.	
12 13 14 15	2.	Analytical Technique: Any analytical technique that meets the following method performance requirements is acceptable.	
16 17	3.	Definitions:	
18 19 20 21		Alcohol by volume (%ABV) A standard measure of how much alcohol (ethanol) is contained in a given volume of an alcoholic beverage (expressed as a volume percent).	
22 23 24		Ethanol The 2-carbon alcohol with a molecular formula of CH ₃ CH ₂ OH. CAS Registry Number: 64-17-5	
25 26 27 28 29		Kombucha Kombucha is a fermented, effervescent tea beverage made by adding a symbiotic culture of bacteria and yeast (SCOBY) to a solution of tea and sugar, and may include other ingredients.	
30 31 32 33		Limit of Quantitation (LOQ) The minimum concentration which quantitative results may be obtained with 95% confidence.	
34 35 36 37 38		Repeatability Variation arising when all efforts are made to keep conditions constant by using the same instrument and operator and repeating during a short time period. Expressed as the repeatability standard deviation (SD _r); or % repeatability relative standard deviation (%RSD _r).	
39 40 41 42 43		Reproducibility The standard deviation or relative standard deviation calculated from among-laboratory data. Expressed as the reproducibility standard deviation (SD _R); or % reproducibility relative standard deviation (% RSD _R).	
44 45 46 47 48		Recovery Factor The fraction or percentage of the analyte that is recovered when the test sample is analyzed using the entire method.	

51 4. Method Performance Requirements:

Analytical range (% ABV)	0.1 to 2.8
Limit of Quantitation (LOQ) (% ABV)	≤ 0.05
Accuracy (% of mean spiked recovery over the range of the assay)	97 to 102
Repeatability (RSD _r) %	≤ 4
Reproducibility (RSD _R)%	≤ 6
ABV= alcohol by volume	

52 53

55

56

57

59 60

61

62

65

67

68

50

54 5. System suitability tests and/or analytical quality control:

Suitable methods will include blank check samples, and check standards at the lowest point and midrange point of the analytical range.

58 6. Reference Material(s):

NIST Standard Reference M	laterial
---------------------------	----------

- 2893 Ethanol-water solution (nom. 0.08 %)2894 Ethanol-water solution (nom. 0.1 %)
 - 2895 Ethanol-water solution (nom. 0.2 %)
- 63 2896 Ethanol-water solution (nom. 0.3 %)
- 64 2897 Ethanol-water solution (nom. 2 %)

66 Sigma Aldrich

459836 200 proof, anhydrous, ≥99.5% (Sigma-Aldrich)

69 Cerilliant CRMs

E-037 Ethanol-80 (5 ampoule multi-pack), 80 mg/dL E-038 Ethanol-100 (5 ampoule multi-pack), 100 mg/dL E-039 Ethanol-200 (5 ampoule multi-pack), 200 mg/dL E-041 Ethanol-150 (10 ampoule multi-pack), 150 mg/dL

E-044 Ethanol-400 (5 ampoule multi-pack), 400 mg/dL

73

74

75

- 71 72
 - LGC Standards
 - BCR-651, beer at 0.505 % (v/v) ethanol.
 - BCR-652, beer at 0.051 % (v/v) ethanol .
- Refer to Annex F: Development and Use of In-House Reference Materials in <u>Appendix F</u>:
 Guidelines for Standard Method Performance Requirements, 19th Edition of the AOAC
- 78 INTERNATIONAL Official Methods of Analysis (2012). Available at:
- 79 http://www.eoma.aoac.org/app_f.pdf
- 80
- 81

82 **7.** Validation Guidance:

- 83
- 84 Appendix F: Guidelines for Standard Method Performance Requirements; 19th Edition of the
- 85 AOAC INTERNATIONAL Official Methods of Analysis (2012). Available at:
- 86 http://www.eoma.aoac.org/app_f.pdf 87
- 88 Appendix K: Guidelines for Dietary Supplements and Botanicals; 19th Edition of the AOAC
- 89 INTERNATIONAL Official Methods of Analysis (2012). Available on line at:
- 90 <u>http://www.eoma.aoac.org/app_k.pdf</u>
- 91
- 92
- 93 8. Maximum Time-To-Result: None