

AOAC INTERNATIONAL

Stakeholder Panel on Agent Detection Assays
(SPADA)

Stakeholder Panel Meeting

March 22 - 23, 2016



2275 Research Blvd.
Conference Room 110
Rockville, MD, 20850
USA



spada@aoac.org

AOAC INTERNATIONAL

Stakeholder Panel on Agent Detection Assays
Stakeholder Panel Meeting
March 22 - 23, 2016



**2275 Research Blvd.
Conference Room 110
Rockville, MD, 20850
USA**



spada@aoac.org

CHAIR BIOGRAPHIES



**LINDA C. BECK, PhD, MT (ASCP)
LEAD SCIENTIST/MICROBIOLOGIST, CBR OFFICE
NAVAL SURFACE WARFARE CENTER**

Co-Chair, AOAC Stakeholder Panel on Agent Detection Assays

Dr. Linda Beck works for the Department of Defense at the Naval Surface Warfare Center Dahlgren Division (NSWCDD) as a Lead Scientist/Microbiologist in the CBR Defense Division. Linda serves as the Navy Chem Bio Rad Nuclear (CBRN) Action Officer in the CBRN Defense T&E Navy Executive Policy Office. Her responsibilities include working on the joint service CBRN Test & Evaluation Capabilities and Methodology effort chaired by the Deputy Under Secretary of the Army, Test and Evaluation (DUSA-T&E).

Prior to her current position, she worked for the Department of Homeland Security (DHS) for three years, and served as the Deputy Program Manager and Director for Laboratory Operations for the BioWatch Program, the biosurveillance system designed to detect select aerosolized biological agents. As Deputy, she provided technical oversight, guidance, and management of the BioWatch Program's daily laboratory operations, National Security Special Events, and surge capability.

Preceding her DHS position, Dr. Beck worked at the NSWCDD and developed and implemented the BioWatch Quality Assurance Samples laboratory, and served as the Program Manager for the DHS effort at Dahlgren. During that tenure, she also served as the Head of the Micro/Molecular Biology Section, supported the development of methods for testing the efficacy of decontaminants on biotoxins, and served as a Chem/Bio Subject Matter Expert on the Hazard Mitigation, Materiel and Equipment Restoration Advance Technology Demonstration program sponsored by the Defense Threat Reduction Agency, Joint Science and Technology Office (DTRA JSTO).

In addition to her Federal government work, Dr. Beck has 15 years of experience in a career in academia. She was a professor in the Biological Sciences Department at the University of Mary Washington prior to her appointment as a professor in the School of Allied Health Professions at the Medical College of Virginia/Virginia Commonwealth University. During her academic tenure, she mentored numerous undergraduate and graduate students through her research in the areas of genetics, microbiology, and cellular biology.

Dr. Beck graduated from the Medical College of Virginia, Virginia Commonwealth University (MCV/VCU) with a PhD in Pathology/Clinical Microbiology followed by two years as a Postdoctoral Research Fellow in the School of Medicine at MCV/VCU.

MATTHEW DAVENPORT, PhD
PROGRAM MANAGER, BIOSCIENCES AND INFORMATICS
THE JOHNS HOPKINS UNIVERSITY APPLIED PHYSICS LABORATORY

Co-Chair, AOAC Stakeholder Panel on Agent Detection Assays

Matt is a Program Manager in Biosciences and Informatics at the Johns Hopkins University Applied Physics Laboratory (JHU/APL) to include projects in personalized genomics, the Microbiome, and functional biology. Matt also works in the areas of human performance and austere medicine with military communities. Prior to JHU/APL, Matt was a Program Manager in the Department of Homeland Security Science and Technology Directorate (DHS S&T) where he established the DHS Public Safety Actionable Assay (PSAA) program and the Stakeholder Panel for Agent Detection Assays (SPADA) to develop voluntary consensus standards for the validation of biothreat detection technologies used by first responders and private-sector end users. In addition to the PSAA program, Matt coordinated a number of bioinformatics efforts including: the development of new databases and software to identify signatures that can be used to specifically detect biothreat agents; sequencing strains of biothreats and their genetic near-neighbors; and application of next generation sequencing to biothreat detection. He also served on numerous interagency committees and co-chaired a working group under the National Science and Technology Council that produced *A National Strategy for CBRNE Standards*.

Matt joined DHS S&T as a Science and Technology Policy Fellow from the American Association for the Advancement of Science (AAAS) where he worked in the same areas of biological countermeasures. Prior to DHS, he was a postdoctoral fellow at both The Johns Hopkins University School of Medicine and the Memorial Sloan-Kettering Cancer Center studying the biochemical mechanisms that control replication of the human genome and the repair of genome when it becomes damaged. Matt earned his doctorate from the Department of Microbiology and Immunology at the University of North Carolina at Chapel Hill and a B.S. in microbiology from North Carolina State University.

Ted Hadfield, PhD**Owner, Hadeco LLC.***SPADA BACILLUS ANTHRACIS WORKING GROUP CO-CHAIR*

Ted L. Hadfield, Ph.D., Co-chair of the Variols Working Group, graduated from University of Utah in 1976. He did a post doctoral in Clinical Immunology at the Latter Day Saints Hospital in Salt Lake City, UT. He subsequently was an assistant professor at California State University in Los Angeles. In 1980 he joined the United States Air Force as a Laboratory Officer. He was stationed at the Armed Forces Institute of Pathology as Chief of Bacteriology. In 1984 he was transferred to Wilford Hall USAF Medical Center in San Antonio Texas as Chief, Clinical Microbiology. In 1989, he transferred back to the Armed Forces Institute of Pathology as Chief of Microbiology. Dr. Hadfield retired from the Air Force in 2000 and was appointed as a Distinguished Scientist at the American Registry of Pathology. He continued as Chief of Microbiology and as Deputy Director of Infectious and Parasitic Diseases Pathology. In 2003 he moved to MRIGlobal's Florida Division as Chief, Bioscience Advisor. In 2012 he retired from MRIGlobal and became president of HADECO, LLC, a consultation service for microbiological, immunology and molecular biology solutions. Dr Hadfield has more than 100 scientific publications and remains active in research projects at MRIGlobal, University of Florida, Gainesville and consultations with clinical laboratories.

Jay E. Gee, PhD**Research Biologist, Bacterial Special Pathogens Branch, DHCPP, NCEZID
United States Centers for Disease Control and Prevention***SPADA BURKHOLDERIA PSEUDOMALLEI WORKING GROUP CHAIR*

Jay E. Gee earned his BS in Microbiology at Mississippi State University in 1987 and his PhD in Biochemistry in 1992 at the University of Alabama at Birmingham School of Medicine. He studied antisense oligonucleotide technology in his first postdoctoral position at Baylor College of Medicine in Houston, TX. He later studied antiviral therapy strategies using chemically modified oligonucleotides in a vesicular stomatitis virus model at L'Institut de Génétique Moléculaire de Montpellier (The Institute of Molecular Genetics of Montpellier) in France in a second postdoctoral position.

He has been with the CDC for almost 14 years. During his research at CDC, he designed real-time PCR assays to identify pathogenic *Leptospira* spp. and *Burkholderia pseudomallei* and has performed molecular genetic subtyping on a variety of pathogens such as *Bacillus* spp. (e.g. *B. anthracis* and *B. cereus*) and *Burkholderia* spp. (e.g. *B. pseudomallei* and *B. mallei*) in support of epidemiological case investigations. He has served on the CDC Environmental Microbiology Work Group and serves on the CDC Next Generation Sequencing Quality Workgroup. He is currently a subject matter expert on *Burkholderia pseudomallei* and *B. mallei*.

Luther Lindler, PhD**Department of Homeland Security, Science and Technology Directorate***SPADA YERSINIA PESTIS WORKING GROUP CHAIR*

Dr. Lindler joined the DHS Science and Technology Directorate in October 2003 as a Senior Science Advisor. Dr. Lindler currently serves as the DHS S&T liaison to the Department of Defense Joint Program Executive Office for Chemical and Biological Defense (JPEO-CBD). He also serves as the Chief Scientist for the DHS Chemical and Biological Defense Division providing biodefense expertise to both DOD as well as DHS in the area of infectious disease threats from a global perspective. Dr. Lindler's previous work provided strategic investments to bring forward deployed rapid molecular diagnostics to U.S. forces. Dr. Lindler provided technical leadership in the Federal Material Threat Assessment and Biological Risk Assessment programs. He helped plan the National Biodefense Analysis and Countermeasures Center forensics and threat characterization programs as well as the first DHS laboratory building on the Fort Detrick National Biodefense Campus. Before joining DHS, Dr. Lindler was a leader in the U.S. Army Biodefense program. He was a principle investigator at the Walter Reed Army Institute of Research leading a team of professionals studying the pathogenesis of the plague bacterium. He served on the Army's plague vaccine steering committee and the emerging threats steering committee within the Biodefense program. The peak of his career with the Army culminated with his senior editorship of the well-acclaimed Biodefense book entitled, "Biological Weapons Defense; Infectious Diseases and Counterbioterrorism." Dr. Lindler was a postdoctoral fellow in the laboratory of Dr. Susan Straley at the University of Kentucky in Lexington from 1987 until 1989. Dr. Lindler received his Ph.D. in Microbiology from the Medical College of Virginia in 1987, his Masters of Science in Microbiology from Clemson University in 1981 and his Bachelor's of Science in Medical Technology from Lenoir Rhyne College in North Carolina in 1978.

Paul Jackson, PhD**Los Alamos and Lawrence Livermore National Laboratories (Retired)***SPADA BACILLUS ANTHRACIS WORKING GROUP CO-CHAIR*

Paul received his Bachelor's of Science degree from the University of Washington in Cellular Biology and his Ph.D. from the University of Utah in Molecular Biology. He was a visiting scholar at the Center for International Security and Cooperation (CISAC) at Stanford University from September 2011-September 2012 and is now a CISAC affiliate. He is also an adjunction professor at the Middlebury Institute of International Studies at Monterey (formerly the Monterey Institute of International Studies) where he team teaches a class entitled "Science and Technology for Non-proliferation and Terrorism Studies".

Paul Jackson (continued)

For the past 24 years he has been studying bacterial pathogens, first working to develop DNA-based methods of detecting these microbes and their remnants in environmental and laboratory samples, then developing methods to differentiate among different strains of the same pathogenic species. Research interests include the study of different methods of interrogating biological samples for detection and characterization of content, and development of bioforensic tools that provide detailed information about biothreat isolates including full interrogation of samples for strain content and other genetic traits. Methods he and collaborators developed have been applied to forensic analysis of samples and aid in identifying the source of disease outbreaks. He contributed to analysis of the *Bacillus anthracis* present in the 2001 Amerithrax letters and conducted detailed analyses of human tissue samples preserved from the 1979 Sverdlovsk anthrax outbreak, providing evidence that was inconsistent with Soviet government claims of a natural anthrax outbreak. His current interests continue to focus on development of assays that rapidly detect specific signatures including antibiotic resistance in threat agents and other pathogens. More recent activities include identification and characterization of new antimicrobial compounds that are based on the pathogens' own genes and the products they encode. These include development of such materials as therapeutic antimicrobials, their application to remediate high value contaminated sites and materials, and their use to destroy large cultures and preparations of different bacterial threat agents. Efforts to address issues of antibiotic resistance and treatment of resistant organisms have recently been expanded to look at non-threat agent pathogens that cause problematic nosocomial or community-acquired infections of particular interest to the military.

Paul spent 24 years as a Technical Staff Member at Los Alamos National Laboratory where he was heavily involved in development of the biological threat reduction efforts there. He was appointed a Laboratory Fellow at Los Alamos – a lifetime appointment - in recognition of his efforts. He moved to Lawrence Livermore National Laboratory in 2005 where he was a Senior Scientist in the Global Security and Physical and Life Sciences Directorates until his retirement in 2013. In addition to his work at the National Laboratories, he has served on the FBI's Scientific Working Group for Microbial Forensics, on NIH study sections and review panels, and continues to serve on steering and oversight committees for other federal agencies.



Frank F. Roberto, PhD, SM (NRCM)
Directorate Fellow, Energy and Environment
Idaho National Laboratory

SPADA BRUCELLA WORKING GROUP CHAIR

Frank Roberto received his BS and PhD in biochemistry from the University of California, Davis, and University of California, Riverside. After a postdoctoral fellowship in molecular plant pathology at UC Davis, he moved to the US Dept. of Energy's Idaho National Laboratory, where he has conducted and directed R&D programs ranging from biomining with acidophilic bacteria and archaea to rapid detection of priority bacterial pathogens such as Brucella. For nearly ten years he worked closely with wildlife biologists studying interspecies transmission of brucellosis to develop field-deployable DNA assays to address bison and elk management issues in the Greater Yellowstone Area. He is a Specialist Microbiologist in biological safety (National Registry of Certified Microbiologists) and has held the Certified Biological Safety Professional (CBSP) certification (American Biological Safety Association).

Shashi Sharma, Ph.D.

SPADA Botulinum neurotoxin A Chair

Dr. Sharma received Ph.D. in Microbiology from University of Bhopal, Bhopal India. After Ph.D., he joined Lupin Biotechnology as a Scientist where he worked on development monoclonal antibodies and immunodiagnosics of HIV, Typhoid and Syphilis. He did posdoc from Department of Biochemistry, University of Massachusetts Dartmouth, where he worked on the structure and function of *Clostridium botulinum* neurotoxin and its associated proteins. Dr. Sharma joined FDA/ CFSAN, in May 2002. His research focuses on the development and validation of an effective and sensitive detection system for *Clostridium botulinum* in foods. He has over 22 years of experience in *C. botulinum* research and published several research papers in peer reviewed journals and holds an US patent on *C. botulinum* toxin associated proteins.

Dr. Victoria Olson
Microbiologist
United States Centers for Disease Control and Prevention

CHAIR, SPADA VARIOLA WORKING GROUP

Victoria Olson obtained her Ph.D. in Biochemistry from the University of Wisconsin – Madison in 2001. Her dissertation focused on understanding transcriptional regulation by the baculovirus *Autographa californica multicausid nucleopolyhedrovirus* immediate early protein (IE1). Dr. Olson then joined the Poxvirus Program at the Centers for Disease Control and Prevention as an Oak Ridge Institute for Science and Education postdoctoral fellow in 2002. Her postdoctoral research focused on understanding how *Orthopoxviruses* interact with their hosts. While studying *Orthopoxviruses*, Dr. Olson completed training and certification for work at multiple biosafety levels, including work with *variola virus* within the Biosafety level 4 laboratories. In 2008, Dr. Olson became lead of the Virus-Host Molecular Interactions Unit within the Poxvirus Team at the Centers for Disease Control and Prevention. She supervises 4 masters-level researchers, 1 post-doctorate, 1 veterinarian, and 1 technician. The Virus-Host Molecular Interactions Unit focuses on research aimed at understanding how *Orthopoxviruses* interact with their hosts and what measures are effective at abrogating disease progression and mitigating morbidity. Since 2005, Dr. Olson has been closely involved in the validation of real-time PCR diagnostic assays for use in clinical settings, with particular focus on obtaining regulatory approvals. During her 12 years within the Poxvirus Team, she has contributed to some 39 peer-reviewed publications.



David Wagner, PhD

Associate Professor, Department of Biological Sciences

Associate Director, Center for Microbial Genetics and Genomics

Northern Arizona University

SPADA F. TULARENSIS WORKING GROUP CO-CHAIR

Dave Wagner has been working with dangerous pathogens, including *Bacillus anthracis*, *Yersinia pestis*, *Francisella tularensis*, and *Burkholderia pseudomallei*, in field and laboratory settings since 1999. He is the Associate Director of the Center for Microbial Genetics and Genomics at NAU, which employs more than 60 faculty, staff, and students. Dr. Wagner has established research collaborations around the world, including *F. tularensis* research in Europe and Asia and *Y. pestis* research in Africa, Asia, Europe, and South America, among many others. He is broadly interested in the evolutionary history, phylogeography, and ecology of infectious disease agents.



The Scientific Association Dedicated to Analytical Excellence®

STAKEHOLDER PANEL ON AGENT DETECTION ASSAYS

Tuesday – Wednesday, March 22-23, 2016

AOAC INTERNATIONAL Headquarters
Conference Room 110
2275 Research Blvd., Rockville, Maryland, 20850
9:00 a.m. – 6:00 p.m.

STAKEHOLDER PANEL AGENDA – MARCH 22, 2016 (Day 1)

- I. Introductions and Call to Order
Jim Bradford, AOAC INTERNATIONAL
- II. Meeting Overview and Objectives (9:05 a.m. – 9:20 a.m.)
Matthew Davenport, DHS, SPADA Co-Chair and Linda Beck, DoD NSWC, SPADA Co-Chair
- III. Discussion on Environmental Factors
Scott Coates, AOAC INTERNATIONAL (9:20 a.m. - 9:35 a.m.)*
- IV. Draft Standard Method Performance Requirements (SMPR) (9:35 a.m. – 1:45 p.m.)
 - a. AOAC Policies and Procedures for Adopting an SMPR – *Deborah McKenzie, AOAC INTL. (9:35 a.m. – 9:40 a.m.)*
 - b. *Bacillus anthracis* - *Paul Jackson, LLNL (ret) and Ted Hadfield, Hadeco LLC * (9:40 a.m. – 10:40 a.m.)*
 - c. *Francisella tularensis* – *Paul Keim, NAU and Dave Wagner, NAU* (11:00 a.m. – 12:00 p.m.)*
 - d. *Yersinia pestis* – *Luther Lindler, DHS* (12:45 p.m. – 1:45 p.m.)*
- V. Launch of New Working Groups (1:45 p.m. – 6:00 p.m.)
 - a. *Variola majora* – *Victoria Olson, CDC (1:45 p.m. – 2:30 p.m.)*
 - i. Fitness for Purpose*
 - b. *Brucella* – *Frank Roberto, Idaho National Lab - (2:30 p.m. – 3:15 p.m.)*
 - i. Fitness for Purpose*
 - c. *Burkholderia pseudomallei*, *Jay Gee, CDC (3:30 p.m. – 4:15 p.m.)*
 - i. Fitness for Purpose*
 - d. *Botulinum neurotoxin A* – *Shashi Sharma, FDA (4:15 p.m. – 5:00 p.m.)*
 - i. Fitness for Purpose*
- VI. Adjourn (5:00 p.m.)

Morning Break: 10:40a.m. – 11:00 p.m.

Lunch: 12:00 – 12:45

Afternoon Break: 3:15pm – 3:30pm

NO GOVERNMENT FUNDS HAVE BEEN USED IN THE PROVISION OF FOOD FOR THIS MEETING

*Item requires a vote

**AOAC INTERNATIONAL Stakeholder Panel on Agent Detection Assays
Working Group Sessions – MARCH 23, 2016 (Day 2)
9:00 a.m. – 4:30 p.m.**

I. AOAC Policies and Procedures for Working Groups (9:00 a.m. – 9:15 a.m.)

Deborah McKenzie, AOAC INTERNATIONAL

II. *Botulinum neurotoxin A* (9:15 a.m. – 10:45 a.m.)

Chair: Shashi Shamra, FDA

- a. Review Fitness for Purpose
SMPR Development

III. *Brucella* (11:00 a.m. – 12:30 p.m.)

Chair: Frank Roberto, Idaho National Lab

- a. Review Fitness for Purpose
- b. SMPR Development

IV. *Burkholderia pseudomallei* (1:30 p.m. – 3:00 p.m.)

Chair: Jay Gee, CDC

- a. Review Fitness for Purpose
- b. SMPR Development

V. *Variola majora* (3:30 p.m. – 5:00 p.m.)

Chair: Victoria Olson, CDC

- a. Review Fitness for Purpose
- b. SMPR Development

Lunch: 12:30 – 1:30

NO GOVERNMENT FUNDS HAVE BEEN USED IN THE PROVISION OF FOOD FOR THIS MEETING

**Item requires a vote*



AOAC Stakeholder Panel on Agent Detection Assays (SPADA)

AOAC STANDARDS DEVELOPMENT PROCESS

APPROVAL OF AN AOAC SMPR®


Deborah McKenzie
Sr. Director,
AOAC Standards Development
AOAC INTERNATIONAL
September 1, 2015

AOAC INTERNATIONAL HEADQUARTERS
2275 Research Blvd., Ste 300
Bethesda, Maryland 20814



The Scientist's Association Dedicated to Analytical Excellence®

AOAC Standards Development Processes




AOAC INTERNATIONAL

As an international standards development organization, AOAC must maintain the following principles throughout all its standard setting activities:

- Transparency
- Openness
- Balance of Interests
- Due Process
- Consensus
- Appeals



Stakeholder Panel Role and Output



AOAC Voluntary Consensus Standards

- **Standard Method Performance Requirements (SMPRs)**
 - Published in *Official Methods of Analysis of AOAC INTERNATIONAL*
 - Manuscript published in *Journal of AOAC INTERNATIONAL*



Stakeholder Panel Composition

- Product Manufacturers
- Analyte/Method Subject Matter Experts
- Technology Providers
- Method Developers
- Government and Regulatory Agencies
- Contract Research Organizations
- Reference Materials Developers
- Ingredient Manufacturers
- Method End Users
- Academia & Research
- Non-Governmental Organizations (ISO, IDF, etc...)
- Other.... as identified

Anyone with a material interest can participate
Balanced group of representative voting stakeholders
Chair and voting members vetted



Organizational Meeting Registrants

- Association of Public Health Laboratories
- ATCC
- bioMérieux
- Censeo Insight
- US Defense Threat Reduction Agency
- FBI
- Hadeco
- Ibis Biosciences
- InSilixa
- Interagency Board
- Gerstel
- JPEO
- Lawrence Livermore National Lab
- Maryland Department of Agriculture
- Minnesota Department of Health
- NIH/NIAID
- Naval Medical Research Center
- New Horizons Diagnostics Corporation
- NIST
- North Carolina DHHS
- Northern Arizona University
- Northrup Grumman
- R-Biopharm
- Texas A&M
- USAMRIID
- US Army Edgewood Chemical Biological Center
- US CDC
- USDA APHIS
- US DHS S&T
- US DHS OHA
- US DOD Critical Reagents Program
- US DoD Navy
- US DoD Dugway Proving Ground
- US FDA-CFSAN
- US FDA ORS




Registered Organizations by Broad Perspectives

- Academia/Research
 - Northern Arizona University
 - Texas A&M
- Industry
 - New Horizons Diagnostics Corporation
 - Northrup Grumman
 - R-Biopharm
 - ATCC
 - bioMérieux
 - Censeo Insight
 - Hadeco
 - Ibis Biosciences
 - InSilixa
 - Gerstel
- Non Governmental Organization
 - APHL
- Government
 - FBI
 - IAB
 - Lawrence Livermore National Lab
 - Maryland Department of Agriculture
 - Minnesota Department of Health
 - NIH/NIAID
 - NIST
 - North Carolina DHHS
 - US CDC
 - USDA APHIS
 - US DHS S&T
 - US DHS OHA
 - US FDA-CFSAN
 - US FDA ORS
- Government, Military
 - JPEO
 - Naval Medical Research Center
 - USAMRIID
 - US Army Edgewood Chemical Biological Center
 - US DOD Critical Reagents Program
 - US DOD Defense Threat Reduction Agency
 - US DoD Navy
 - US DoD Dugway Proving Ground




Registrants by Broad Perspectives




Registrants by Specific Perspectives

- Coordinator
- Method Developer
- Public Health
- Regulatory
- Evaluation
- Method End User
- Public Safety
- Research
- Independent
- Programs
- Reference Materials
- Technology Provider




Registrants by Broad and Specific Perspectives

- Government - Coordinator
- Government - Independent
- Industry - Method Developer
- Industry - End User
- Government - State Public Health
- Government - Reference Materials
- Government - Regulatory
- Academia - Research
- Military - Research
- Military Evaluation
- Industry - Independent
- Military - Method Developer
- Military - Programs
- NGO - Public Health
- Industry - Reference Materials
- Government - State Regulatory
- Government - Research
- Industry - Technology Provider




Registrants by Region - In/Out of USA



Approving AOAC Standards

- Working Group Chair or designee will present on the draft standard method performance requirements including reconciled comments received on behalf of the working group and moves for SPADA to adopt the SMPR® as presented
- SPADA chair will entertain deliberation on the draft standard
- After due deliberation, SPADA chair will call for a vote
- Voting members will be able to vote in favor of the motion, against the motion, or abstain from voting
- 2/3 vote in favor required to approve/adopt an AOAC SMPR®.



Documentation and Communication

- AOAC carefully documents the actions of Stakeholder Panel and the Working Groups
- AOAC will prepare summaries of the meetings
 - Communicate summaries to the stakeholders
 - Publish summaries in the *Referee* section of AOAC's *Inside Laboratory Management*
- AOAC publishes its voluntary consensus standards and *Official Methods*
 - *Official Methods of Analysis of AOAC INTERNATIONAL*
 - *Journal of AOAC INTERNATIONAL*
- AOAC publishes the status of standards in the Referee section of AOAC's *Inside Laboratory Management*



Roles and Responsibilities

- Stakeholder Panel
 - Establish working groups to develop standards
 - Comment, deliberate, and establish voluntary consensus standards
- Stakeholder Panel Working Groups
 - Develop draft standard method performance requirements
 - Reconcile comments
 - Present draft standard to stakeholders
- Official Methods Board
 - Vet and approve stakeholder panel chair and representative voting members
 - Assign representative to serve as a resource to stakeholder panel
- AOAC Staff
 - Coordinate stakeholder panel, working groups, and facilitate their meetings.
 - Document actions/decisions of working groups and stakeholder panel
 - Post SMPRs and collect comments for draft SMPRs



Contact Information


Contact AOAC Staff:

Tel: 301.924.7077

Web: www.aoac.org

- **E. James Bradford**, Executive Director & CEO, jbradford@aoac.org, ext. 102
- **Krystyna McIver**, Executive for Scientific Engagement and Communication, kmciver@aoac.org, ext. 111
- **Scott Coates**, Chief Scientific Officer, scoates@aoac.org, ext. 137
- **Deborah McKenzie**, Sr. Director – Standards Development and AOAC Research Institute, dmckenzie@aoac.org, ext. 157





AOAC INTERNATIONAL STAKEHOLDER PANEL ON AGENT DETECTION ASSAYS

Bacillus anthracis Working Group
Co-Chairs: Ted Hadfield, Hadecco and Paul Jackson, LLNL (ret)
SMPR Presentation
March 22, 2016

Rockville, Maryland, USA

Fitness for Purpose

“Field-deployable detection of *Bacillus anthracis* in samples from aerosol collection devices, using nucleic acid-based techniques developed for the Department of Defense.”



SPADA *Bacillus anthracis* Working Group Members

Ted Hadfield, Hadeco (Co-Chair)	Crystal Jaing, LLNL
Paul Jackson, LLNL (Ret.)	Malcolm Johns, DHS
Jessica Appler, HHS/BARDA	Nancy Lin, NIST
Les Ballie, Cardiff University	Laura Maple, NSWC
Ed Bailor, IAB	Stephen Morse, CDC
Jeff Ballin, ECBC	Dallas New,
Timothy Bauer, NSWC	Michael Retford, JBTS JPM NBCCA
Linda Beck, NSWC	Sanjiv Shah, US EPA
Steven Blanke, University of Illinois	Snahmuga Sozhamannan, CRP
Ryan Cahall, Censeo Insight	David Trudil, New Horizons
Ken Damer, Northrop Grumman	Susan Welkos, USAMRIID



***Bacillus anthracis* Working Group - Work to Date**

- Working Group Launch (September, 2015)
- Four teleconferences (October – December 2015)
- One SMPR Drafted
- Public comment period (January 8, 2016 – February 5, 2016)
- SMPR made ready for SPDS review and approval



SMPR Key Points

Table 1: Methods Performance Requirements

Parameter	Minimum Performance Requirement
AMDL	2,000 standardized BA Ames spores per mL liquid in the candidate method sample collection buffer.
Probability of Detection at AMDL within sample collection buffer	≥ 0.95
Probability of Detection at AMDL in environmental matrix materials.	≥ 0.95
System False-Negative Rate using spiked environmental matrix materials.	≤ 5%
System False-Positive Rate using environmental matrix materials.	≤ 5%
Inclusivity	All inclusivity strains (Table III) must test positive at 2x the AMDL [†]
Exclusivity	All exclusivity strains (Table IV and Table V; part 2) must test negative at 10x the AMDL [†]
Notes: [†] 100% correct analyses are expected. All discrepancies are to be re-tested following the AOAC Guidelines for Validation of Biological Threat Agent Methods and/or Procedures ² .	



² Official Methods of Analysis of AOAC INTERNATIONAL (2012) 19th Ed., AOAC INTERNATIONAL, Gaithersburg, MD, USA, APPENDIX I; also on-line at http://www.eoma.aocac.org/app_1.pdf.

SMPR Key Points

Table II: Controls

Control	Description	Implementation
Positive Control	This control is designed to demonstrate an appropriate test response. The positive control should be included at a low but easily detectable concentration, and should monitor the performance of the entire assay. The purpose of using a low concentration of positive control is to demonstrate that the assay sensitivity is performing at a previously determined level of sensitivity.	Single use per sample (or sample set) run
Negative Control	This control is designed to demonstrate that the assay itself does not produce a detection in the absence of the target organism. The purpose of this control is to rule-out causes of false positives, such as contamination in the assay or test.	Single use per sample (or sample set) run
Inhibition Control	This control is designed to specifically address the impact of a sample or sample matrix on the assay's ability to detect the target organism.	Single use per sample (or sample set) run



SMPR Key Points

Table III: Inclusivity Panel

No.	Cluster	Genotype	Strain	Origin	Characteristics
1	A1a	7	Canadian bison	Wood bison	pXO1 ^a , pXO2 ^a , VNTR ^a genotype group A1a
2	A3a	45 ^b	V770-NP-1R	Vaccine (USA)	pXO1 ^a , pXO2 ^a , VNTR genotype group A3A
3	A2	29	PAK-1	Sheep (Pakistan)	pXO1 ^a , pXO2 ^a , VNTR genotype group A2
4	A3a	51	BA1015	Bovine (MD)	pXO1 ^a , pXO2 ^a , VNTR genotype group A3a
5	A3b	62	Ames	Bovine (Texas)	pXO1 ^a , pXO2 ^a , VNTR genotype group A3b
6	A3c	67	K3	South Africa	pXO1 ^a , pXO2 ^a , VNTR genotype group A3c
7	A3d	68	Ohio ACB	Pig	pXO1 ^a , pXO2 ^a , VNTR genotype group A3d
8	A4	69	SK-102 (Pakistan)	Imported wool	pXO1 ^a , pXO2 ^a , VNTR genotype group A4
9	A4	77	Vollum 1B	USAMRIID ^c	pXO1 ^a , pXO2 ^a , VNTR genotype group A4
10	B1	82	BA1035	Human (S. Africa)	pXO1 ^a , pXO2 ^a , VNTR genotype group B1
11	B2	80	RA3	Bovine (France)	pXO1 ^a , pXO2 ^a , VNTR genotype group B2
12	A1a	8	Pasteur	USAMRIID ^c	pXO1 ^a , pXO2 ^a , VNTR genotype group A1a
13	A3b	59, 61 ^b	Sterne	USAMRIID ^c	pXO1 ^a , pXO2 ^a , VNTR genotype group A3b
14	A1b	23	Turkey No. 32	Human (Turkey)	pXO1 ^a , pXO2 ^a , VNTR genotype group A1b

^a VNTR: Variable number tandem repeat

^b Organism contains only seven of eight multiple locus variable number tandem repeat analysis (MLVA) markers due to the absence of pXO2. Genotypes listed are consistent with seven of the eight markers.

^c USAMRIID = The United States Army Medical Research Institute for Infectious Diseases.



Background


Inclusivity panel: Strains picked and rationale for selection

- All inclusivity panel isolates must meet clinical criteria for being *B. anthracis*
 - Colony morphology, Gram stain, presence of capsule, motility, hemolytic activity, γ phage sensitivity, pen^s, etc.
 - An isolate must contain the plasmids pXO1 and pXO2 unless otherwise identified as missing one of these plasmids
 - An isolate that maps, by multiple phylogenetic methods, to a tight cluster of microbes sharing these physical and pathogenic characteristics




Background

Inclusivity panel: Strains picked and rationale for selection



Background

Inclusivity panel: Strains picked and rationale for selection



From Keim, *et al.* (2000)
J. Bacteriol. 182, 2928-2936



SMPR Key Points

Table IV: Exclusivity Panel (near-neighbor)

No.	Species	Strain	Plasmid status
1	<i>B. cereus</i>	S2-8	pXO1 ^a , pXO2
2	<i>B. cereus</i>	3A	pXO1 ^a , pXO2
3	<i>B. thuringiensis</i>	HD1011	pXO1 ^a , pXO2
4	<i>B. thuringiensis</i>	HD682	pXO1 ^a , pXO2
5	<i>B. cereus</i>	D17	pXO1 ^a , pXO2
6	<i>B. thuringiensis</i>	HD571	pXO1 ^a , pXO2
7	<i>B. cereus</i>	Al Hakam	pXO1 ^a , pXO2
8	<i>B. cereus</i>	ATCC 4342	pXO1 ^a , pXO2
9	<i>B. cereus</i>	FM1	pXO1 ^a , pXO2
10	<i>B. cereus</i>	E33L	pXO1 ^a , pXO2
11	<i>B. thuringiensis</i>	97-27	pXO1 ^a , pXO2
12	<i>B. cereus</i>	G9241	pBCXO1 ^a , pXO2
13	<i>B. cereus</i>	03BB102	pXO1 ^a , capA ⁺ , capB ⁺ , capC ^{1b}
14	<i>B. cereus</i>	03BB108	pXO1 ^a , capA ⁺ , capB ⁺ , capC ^{1b}
15	<i>B. cereus</i> subsp. <i>anthracis</i>		

This isolate was eliminated from the panel



^a pBCXO1 is pXO1-like, but not identical.
^b capA, capB, and capC are contained within the *Bacillus anthracis* pXO2 plasmid; however, the capA, capB, and capC sequences are found in strains 03BB102 and 03BB108 in the absence of the pXO2 plasmid.



Guidance on Combining DNA for Exclusivity Evaluation
 DNA from exclusivity panel organisms 1 -9 in Table IV may be tested as isolated DNA, or combined to form a pool of exclusivity panel organisms, with each panel organism represented at 10 times the AMDL. If an unexpected result occurs, each of the exclusivity organisms from a failed pool must be individually re-tested at 10 times the AMDL.
 DNA from exclusivity panel organisms 10 – 15 in Table IV **can NOT** be combined for exclusivity evaluation.



Background

**Exclusivity panel:
 Strains picked and rationale for selection**






Background

Exclusivity panel: Strains picked and rationale for selection

B. thuringiensis isolates used in U.S. EPA-approved pesticides

- *Bacillus thuringiensis* (Berliner) (no subsp. given)
- *Bacillus thuringiensis* subsp. *aizawai*
- *Bacillus thuringiensis* subsp. *israelensis*
- *Bacillus thuringiensis* subsp. *kurstaki*
- *Bacillus thuringiensis* subsp. *tenebrionis**

*aka subsp. *morrisoni*



Background

Exclusivity panel: Strains picked and rationale for selection

The last three isolates in the Exclusivity Panel do not map closely to *B. anthracis*



SMPR Key Points

Table V: Environmental Factors For Validating Biological Threat Agent Detection Assays
[Adapted from the Environmental Factors Panel approved by SPADA on June 10, 2010.]

The Environmental Factors Studies supplement the biological threat agent near-neighbor exclusivity testing panel. There are three parts to Environmental Factors studies: part 1 – environmental matrix samples; part 2 - the environmental organisms study; and part 3 – the potential Interferents applicable to Department of Defense applications.³

Part 1:

Environmental Matrix Samples - Aerosol Environmental Matrices

Method developers shall obtain environmental matrix samples that are representative and consistent with the collection method that is anticipated to ultimately be used in the field. This includes considerations that may be encountered when the collection system is deployed operationally such as collection medium, duration of collection, diversity of geographical areas that will be sampled, climatic/environmental conditions that may be encountered and seasonal changes in the regions of deployment.

Justifications for the selected conditions that were used to generate the environmental matrix and limitations of the validation based on those criteria must be documented.

- Method developers shall test the environmental matrix samples for interference using samples inoculated with a target biological threat agent sufficient to achieve 95% probability of detection.
- Cross-reactivity testing will include sufficient samples and replicates to ensure each environmental condition is adequately represented.

³Added in June 2015 for the Department of Defense project



SMPR Key Points

Part 2: Environmental Panel Organisms - This list is comprised of identified organisms from the environment.

Inclusion of all environmental panel organisms is not a requirement if a method developer provides appropriate justification that the intended use of the assay permits the exclusion of specific panel organisms. Justification for exclusion of any environmental panel organism(s) must be documented and submitted.

Organisms and cell lines may be tested as isolated DNA, or as pools of isolated DNA. Isolated DNA may be combined into pools of up to 10 panel organisms, with each panel organism represented at 10 times the AMDL, where possible. The combined DNA pools are tested in the presence (at 2 times the AMDL) and absence of the target gene or gene fragment. If an unexpected result occurs, each of the individual environmental organisms from a failed pool must be individually re-tested at 10 times the AMDL with and without the target gene or gene fragment at 2x the AMDL in the candidate method DNA elution buffer.

DNA in this list that already appear in the inclusivity or exclusivity panel do not need to be tested again as part of the environmental factors panel.

- **Potential bacterial biothreat agents**

- Bacillus anthracis* Ames
 - Yersinia pestis* Colorado-92
 - Francisella tularensis* subsp. *tularensis* Schu-S4
 - Burkholderia pseudomallei*
 - Burkholderia mallei*
 - Brucella melitensis*



SMPR Key Points

Cultivable bacteria identified as being present in air soil or water

<i>Acinetobacter lwoffii</i>	<i>Fusobacterium nucleatum</i>	Microbial eukaryotes
<i>Agrobacterium tumefaciens</i>	<i>Lactobacillus plantarum</i>	<u>Freshwater amoebae</u>
<i>Bacillus amyloliquefaciens</i>	<i>Legionella pneumophila</i>	<i>Acanthamoeba castellanii</i>
<i>Bacillus cohnii</i>	<i>Listeria monocytogenes</i>	<i>Naegleria fowleri</i>
<i>Bacillus psychrosaccharolyticus</i>	<i>Moraxella nonliquefaciens</i>	
<i>Bacillus benzoovorans</i>	<i>Mycobacterium smegmatis</i>	<u>Fungi</u>
<i>Bacillus megaterium</i>	<i>Neisseria lactamica</i>	<i>Alternaria alternata</i>
<i>Bacillus horikoshii</i>	<i>Pseudomonas aeruginosa</i>	<i>Aspergillus fumigatus</i>
<i>Bacillus macroides</i>	<i>Rhodobacter sphaeroides</i>	<i>Aureobasidium pullulans</i>
<i>Bacteroides fragilis</i>	<i>Riemerella anatipestifer</i>	<i>Cladosporium cladosporioides</i>
<i>Burkholderia cepacia</i>	<i>Shewanella oneidensis</i>	<i>Cladosporium sphaerospermum</i>
<i>Burkholderia gladioli</i>	<i>Staphylococcus aureus</i>	<i>Epicoccum nigrum</i>
<i>Burkholderia stabilis</i>	<i>Stenotrophomonas maltophilia</i>	<i>Eurotium amstelodami</i>
<i>Burkholderia plantarii</i>	<i>Streptococcus pneumoniae</i>	<i>Mucor racemosus</i>
<i>Chryseobacterium indologenes</i>	<i>Streptomyces coelicolor</i>	<i>Paecilomyces variotii</i>
<i>Clostridium sardiniense</i>	<i>Synechocystis</i>	<i>Penicillium chrysogenum</i>
<i>Clostridium perfringens</i>	<i>Vibrio cholerae</i>	<i>Walleimia sebi</i>
<i>Deinococcus radiodurans</i>		
<i>Delftia acidovorans</i>		
<i>Escherichia coli</i> K12		



SMPR Key Points

DNA from higher eukaryotes

Plant Pollen⁴

Zea mays (corn)
Pinus spp. (pine)
Gossypium spp. (Cotton)

Arthropods

Aedes aegypti (ATCC /CCL-125(tm) mosquito cell line)
Aedes albopictus (Mosquito C6/36 cell line)
Dermatophagoides pteronyssinus (Dust mite -commercial source)
Xenopsylla cheopis Flea (Rocky Mountain labs)
Drosophila cell line
Musca domestica (housefly) ARS, USDA, Fargo, ND
Gypsy moth cell lines LED652Y cell line (baculovirus)- Invitrogen
Cockroach (commercial source)
Tick (Amblyomma and Dermacentor tick species for *F. tularensis* detection assays)⁵

Vertebrates

Mus musculus (ATCC/IB-123) mouse
Rattus norvegicus (ATCC/CRL-1896) rat
Canis familiaris(ATCC/CCL-183) dog
Felis catus (ATCC/CR-8727) cat
Homo sapiens (HeLa cell line ATCC/CCL-2) human
Gallus gallus domesticus (Chicken)
Capra hircus (Goat)⁶

⁴ If pollen is unavailable, vegetative DNA is acceptable

⁵ Added by SPADA on (future approval date).

⁶ Added by SPADA on September 1, 2015



SMPR Key Points

Biological insecticides – Strains of *B. thuringiensis* present in commercially available insecticides have been extensively used in hoaxes and are likely to be harvested in air collectors. For these reasons, it should be used to assess the specificity of these threat assays.

B. thuringiensis subsp. *israelensis*⁷
B. thuringiensis subsp. *kurstaki*⁷
B. thuringiensis subsp. *morrisoni*⁷
Serenade (Fungicide) *B. subtilis* (QST713)

Viral agents have also been used for insect control. Two representative products are:

- Gypcheck for gypsy moths (*Lymantria dispar* nuclear polyhedrosis virus)
- Cyd-X for codling moths (Codling moth granulosis virus)

⁷There are part of the exclusivity panel for testing *B. anthracis*



Background

Johnson, S.L., Daligault, H.E., Davenport, K.W., Jaissle, J., Grey, K.G., Ladner, J.T., Broomall, S.M., Bishop-Lilly, K.A., Bruce, D.C., Gibbons, H.S., Coyne, S.R., Lo, C.-C., Meincke, I., Munk, A.C., Koroleva, G.I., Rosenzweig, C.N., Palacios, f., Redden, C.L., Minogue, T.D. and Chain, P.S. (2015)

Complete Genome Sequences of 35 Biothreat Assay-Relevant *Bacillus* Species.

Genome Announc. 3, e00151-15.

ABSTRACT: In 2011, the Association of Analytical Communities (AOAC) International released a list of *Bacillus* strains relevant to biothreat molecular detection assays. We present the complete and annotated genome assemblies of the 15 strains listed on the inclusivity panel, as well as the 20 strains listed on the exclusivity panel.



Background

- Given full genome sequences for all members of the Inclusivity and Exclusivity panels, anyone designing assays using DNA sequences can use *in silico* methods to select sequences to target with their assays that are specific to *B. anthracis*, based on signatures found only in genomes of isolates in the Inclusivity panel but absent from genomes of organisms in the Exclusivity panel.
- It will be straightforward to demonstrate that the assays were designed in such a manner and, *in silico* "testing" will quickly demonstrate whether the targeted sequences – or sets of sequences – are specific to the target.

Isolates developed using such methods must still be tested against the inclusivity, exclusivity and environmental factors panels to validate the assays because the DNA databases used to develop such assays are still incomplete and do not represent the breadth of isolates found in the environment.



SMPR Key Points

Part 3: Potential Interferents Study

The Potential Interferents Study supplements the Environmental Factors Study, and is applicable to all biological threat agent detection assays for Department of Defense applications. Table VI provides a list of potential Interferents that are likely to be encountered in various Department of Defense applications.

Method developers and evaluators shall determine the most appropriate potential Interferents for their application. Interferents shall be spiked at a final test concentration of 1 µg/ml directly into the sample collection buffer. Sample collection buffers spiked with potential Interferents shall be inoculated at 2 times the AMDL (or AMIL) with one of the target biological threat agents.

Spiked / inoculated sample collection buffers shall be tested using the procedure specified by the candidate method. A candidate method that fails at the 1 microgram per ml level may be reevaluated at lower concentrations until the inhibition level is determined.

It is expected that all samples are correctly identified as positive.



SMPR Key Points

Table VI: Potential Interferents

Compounds	Potential Theaters of Operation	
Group 1: petroleum-based	JP-8 ¹	Airfield
	JP-5 ²	Naval
	diesel/gasoline mixture	Ground
	fog oil (standard grade fuel number 2)	Naval, Ground
	burning rubber ³	Ground, Airfield
Group 2: exhaust	gasoline exhaust	Ground
	jet exhaust	Naval, Airfield
	diesel exhaust	Ground
Group 3: obscurants	terephthalic acid ⁴	Ground
	zinc chloride smoke ⁵	Ground
	solvent yellow ⁶	Ground
Group 4: environmental	burning vegetation	ground, airfield
	road dust	Ground
	sea water (sea spray)	Naval
Group 5: chemicals	brake fluid ⁷	All
	brake dust ⁸	Ground
	cleaning solvent, MIL-L-63460 ⁹	All
	explosive residues a) high explosives ¹⁰ b) artillery propellant ¹¹	All

Table VI is offered for guidance and there are no mandatory minimum requirements for the number of potential Interferents to be tested.




SMPR Key Points

¹**JP-8.** Air Force formulation jet fuel.

²**JP-5.** A yellow kerosene-based jet fuel with a lower flash point developed for use in aircraft stationed aboard aircraft carriers, where the risk from fire is particularly great. JP-5 is a complex mixture of hydrocarbons, containing alkanes, naphthenes, and aromatic hydrocarbons.


³**Burning rubber** (tire smoke). Gaseous C1-C5 hydrocarbons: methane; ethane; isopropene; butadiene; propane. Polycyclic aromatic hydrocarbons (58-6800 ng/m³): parabenzo(a)pyrene; polychlorinated dibenzo-p-dioxins (PCDD); polychlorinated dibenzofurans (PCDF). Metals (0.7 - 8 mg/m³): zinc; lead; cadmium.

⁴**Terephthalic acid.** Used in the AN/M83 hand grenade currently used by US military.



⁵**Zinc chloride smoke.** Also known as "zinc chloride smoke" and "HC smoke". Was used in the M8 grenade and still used in 155mm artillery shells. HC smoke is composed of 45% hexachloroethane, 45% zinc oxide, and 10% aluminum.

⁶**Solvent yellow 33** [IUPAC name: 2-(2-quinoly)-1,3-indandione] is a new formulation being developed for the M18 grenade.



SMPR Key Points

⁷**Brake fluid.** DOT 4 is primarily composed of glycol and borate esters. DOT 5 is silicone-based brake fluid. The main difference is that DOT 4 is hygroscopic whereas DOT 5 is hydrophobic. DOT 5 is often used in military vehicles because it is more stable over time requires less maintenance.

⁸**Brake dust.** Fe particles caused by abrasion of the cast iron brake rotor by the pad and secondly fibers from the semi metallic elements of the brake pad. The remainder of the dust residue is carbon content within the brake pad.

⁹**MIL-L-63460,** "Military Specification, Lubricant, Cleaner and Preservative for Weapons and Weapons Systems"; trade name "Break-Free CLP". Hyperlink: [Midway USA](#).

¹⁰**High explosives.** The M795 155mm projectile is the US Army / Marine Corp's current standard projectile containing 10.8 kg of TNT. The M795 projectile replaced the M107 projectile that contained Composition B which is a 60/40 mixture of RDX/TNT. RDX is cyclotrimethylene trinitramine. Suggestion: test RDX/TNT together.

¹¹**Artillery propellant.** Modern gun propellants are divided into three classes: single-base propellants which are mainly or entirely nitrocellulose based, double-base propellants composed of a combination of nitrocellulose and nitroglycerin, and triple base composed of a combination of nitrocellulose and nitroglycerin and nitroguanidine. Suggestion: test total nitrocellulose/ nitroglycerin nitroguanidine together.



Comments Submitted (if any)



Motion

- Motion to accept the Standard Method Performance Requirements for *Bacillus anthracis* as presented.



Discussion?



4 **Standard Method Performance Requirements (SMPRs®) for**
5 **DNA-based methods of detecting *Bacillus anthracis* in field-deployable, Department of**
6 **Defense aerosol collection devices**
7

8 **Intended Use:** *Field-deployed use for analysis of aerosol collection filters and/or liquids*
9

10 **1. Applicability:** Detection of *Bacillus anthracis* in collection buffers from aerosol
11 collection devices. Field-deployable assays are preferred.
12

13 **2. Analytical Technique:** Molecular detection of nucleic acid.
14

15 **3. Definitions:**
16

17 **Acceptable Minimum Detection Level (AMD L)**

18 The predetermined minimum level of an analyte, as specified by an expert committee which
19 must be detected by the candidate method at a specified probability of detection (POD).
20

21 **Environmental Factors**

22 For the purposes of this SMPR: any factor in the operating environment of an analytical
23 method, whether abiotic or biotic, that might influence the results of the method.
24

25 **Exclusivity**

26 Study involving pure non-target strains, which are potentially cross-reactive, that shall not
27 be detected or enumerated by the candidate method.
28

29 **Inclusivity**

30 Study involving pure target strains that shall be detected or enumerated by the candidate
31 method.
32

33 **Interferents**

34 A . . . substance in analytical procedures . . . that, at a (the) given concentration, causes a
35 systematic error in the analytical result.¹ Sometimes also known as interferants.
36

37 **Maximum Time-To- Result**

38 Maximum time to complete an analysis starting from the collection buffer to assay result.
39

40 **Probability of Detection (POD)**

41 The proportion of positive analytical outcomes for a qualitative method for a given matrix at
42 a specified analyte level or concentration with a ≥ 0.95 confidence interval.
43

¹ International Union Of Pure And Applied Chemistry Analytical Chemistry Division Commission On Analytical Reactions And Reagents* Definition And Classification Of Interferences In Analytical Procedures Prepared For Publication By W. E. Van Der Linden. Pure & Appl. Chem., Vol. 61, No. 1, pp. 91-95, 1989. Printed in Great Britain. @ 1989 IUPAC

44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77

System False Negative Rate

Proportion of test results that are negative contained within a population of known positives

System False Positive Rate

Proportion of test results that are positive contained within a population of known negatives.

4. Method Performance Requirements:

See Table I.

5. System Suitability Tests and/or Analytical Quality Control:

The controls listed in Table II shall be embedded in assays as appropriate. Manufacturer must provide written justification if controls are not embedded in the assay.

6. Validation Guidance: AOAC INTERNATIONAL Methods Committee Guidelines for Validation of Biological Threat Agent Methods and/or Procedures (AOAC INTERNATIONAL Official Methods of Analysis, 2012, Appendix I).

Inclusivity and exclusivity panel organisms used for evaluation must be characterized and documented to truly be the species and strains they are purported to be.

8. Maximum Time-to-Result: Within four hours.

78
79

Table I: Method Performance Requirements

Parameter	Minimum Performance Requirement
AMDL	2,000 standardized BA Ames spores per mL liquid in the candidate method sample collection buffer.
Probability of Detection at AMDL within sample collection buffer	≥ 0.95
Probability of Detection at AMDL in environmental matrix materials.	≥ 0.95
System False-Negative Rate using spiked environmental matrix materials.	$\leq 5\%$
System False-Positive Rate using environmental matrix materials.	$\leq 5\%$
Inclusivity	All inclusivity strains (Table III) must test positive at 2x the AMDL [†]
Exclusivity	All exclusivity strains (Table IV and Table V; part 2) must test negative at 10x the AMDL [†]
Notes: [†] 100% correct analyses are expected. All discrepancies are to be re-tested following the AOAC Guidelines for Validation of Biological Threat Agent Methods and/or Procedures ² .	

80
81

² Official Methods of Analysis of AOAC INTERNATIONAL (2012) 19th Ed., AOAC INTERNATIONAL, Gaithersburg, MD, USA, APPENDIX I; also on-line at http://www.eoma.aoac.org/app_i.pdf.

82 **TABLE II: Controls**
 83

Control	Description	Implementation
Positive Control	<p>This control is designed to demonstrate an appropriate test response. The positive control should be included at a low but easily detectable concentration, and should monitor the performance of the entire assay. The purpose of using a low concentration of positive control is to demonstrate that the assay sensitivity is performing at a previously determined level of sensitivity.</p>	<p>Single use per sample (or sample set) run</p>
Negative Control	<p>This control is designed to demonstrate that the assay itself does not produce a detection in the absence of the target organism. The purpose of this control is to rule-out causes of false positives, such as contamination in the assay or test.</p>	<p>Single use per sample (or sample set) run</p>
Inhibition Control	<p>This control is designed to specifically address the impact of a sample or sample matrix on the assay's ability to detect the target organism.</p>	<p>Single use per sample (or sample set) run</p>

84
85
86
87

Table III: Inclusivity Panel

No.	Cluster	Genotype	Strain	Origin	Characteristics
1	A1a	7	Canadian bison	Wood bison	pXO1 ⁺ , pXO2 ⁺ , VNTR ^a genotype group A1a
2	A3a	45 ^b	V770-NP-1R	Vaccine (USA)	pXO1 ⁺ , pXO2 ⁻ , VNTR genotype group A3A
3	A2	29	PAK-1	Sheep (Pakistan)	pXO1 ⁺ , pXO2 ⁺ , VNTR genotype group A2
4	A3a	51	BA1015	Bovine (MD)	pXO1 ⁺ , pXO2 ⁺ , VNTR genotype group A3a
5	A3b	62	Ames	Bovine (Texas)	pXO1 ⁺ , pXO2 ⁺ , VNTR genotype group A3b
6	A3c	67	K3	South Africa	pXO1 ⁺ , pXO2 ⁺ , VNTR genotype group A3c
7	A3d	68	Ohio ACB	Pig	pXO1 ⁺ , pXO2 ⁺ , VNTR genotype group A3d
8	A4	69	SK-102 (Pakistan)	Imported wool	pXO1 ⁺ , pXO2 ⁺ , VNTR genotype group A4
9	A4	77	Vollum 1B	USAMRIID ^c	pXO1 ⁺ , pXO2 ⁺ , VNTR genotype group A4
10	B1	82	BA1035	Human (S. Africa)	pXO1 ⁺ , pXO2 ⁺ , VNTR genotype group B1
11	B2	80	RA3	Bovine (France)	pXO1 ⁺ , pXO2 ⁺ , VNTR genotype group B2
12	A1a	8	Pasteur	USAMRIID	pXO1 ⁻ , pXO2 ⁺ , VNTR genotype group A1a
13	A3b	59, 61 ^b	Sterne	USAMRIID	pXO1 ⁺ , pXO2 ⁻ , VNTR genotype group A3b
14	A1b	23	Turkey No. 32	Human (Turkey)	pXO1 ⁺ , pXO2 ⁺ , VNTR genotype group A1b

88 ^a VNTR: Variable number tandem repeat
89 ^b Organism contains only seven of eight multiple locus variable number tandem repeat analysis (MLVA)
90 markers due to the absence of pXO2. Genotypes listed are consistent with seven of the eight
91 markers.
92 ^c USAMRIID = The United States Army Medical Research Institute for Infectious Diseases.
93
94

95 **Table IV: Exclusivity Panel (near-neighbor)**

No.	Species	Strain	Plasmid status
1	<i>B. cereus</i>	S2-8	pXO1 ⁻ , pXO2 ⁻
2	<i>B. cereus</i>	3A	pXO1 ⁻ , pXO2 ⁻
3	<i>B. thuringiensis</i>	HD1011	pXO1 ⁻ , pXO2 ⁻
4	<i>B. thuringiensis</i>	HD682	pXO1 ⁻ , pXO2 ⁻
5	<i>B. cereus</i>	D17	pXO1 ⁻ , pXO2 ⁻
6	<i>B. thuringiensis</i>	HD571	pXO1 ⁻ , pXO2 ⁻
7	<i>B. cereus</i>	Al Hakam	pXO1 ⁻ , pXO2 ⁻
8	<i>B. cereus</i>	ATCC 4342	pXO1 ⁻ , pXO2 ⁻
9	<i>B. cereus</i>	FM1	pXO1 ⁻ , pXO2 ⁻
10	<i>B. cereus</i>	E33L	pXO1 ⁻ , pXO2 ⁻
11	<i>B. thuringiensis</i>	97-27	pXO1 ⁻ , pXO2 ⁻
12	<i>B. cereus</i>	G9241	pBCXO1 ^{+a} , pXO2 ⁻
13	<i>B. cereus</i>	03BB102	pXO1 ⁺ , capA ⁺ , capB ⁺ , capC ^{+b}
14	<i>B. cereus</i>	03BB108	pXO1 ⁺ , capA ⁺ , capB ⁺ , capC ^{+b}
15	<i>B. cereus</i> subsp. <i>anthracis</i>		

96

97 ^a pBCXO1 is pXO1-like, but not identical.98 ^b capA, capB, and capC are contained within the *Bacillus anthracis* pXO2 plasmid; however, the capA,
99 capB, and capC sequences are found in strains 03BB102 and 03BB108 in the absence of the pXO2
100 plasmid.

101

102

103

Guidance on Combining DNA for Exclusivity Evaluation

104

DNA from exclusivity panel organisms 1 -9 in Table IV may be tested as isolated DNA, or
105 combined to form a pool of exclusivity panel organisms, with each panel organism represented
106 at 10 times the AMDL. If an unexpected result occurs, each of the exclusivity organisms from a
107 failed pool must be individually re-tested at 10 times the AMDL.

108

109

DNA from exclusivity panel organisms 10 – 15 in Table IV can NOT be combined for exclusivity
110 evaluation.

111

112

113

114

115

116 **Table V: Environmental Factors For Validating Biological Threat Agent Detection**
117 **Assays**

118

119 [Adapted from the Environmental Factors Panel approved by SPADA on June 10, 2010.]

120

121 The Environmental Factors Studies supplement the biological threat agent near-neighbor
122 exclusivity testing panel. There are three parts to Environmental Factors studies: part 1 -
123 environmental matrix samples; part 2 - the environmental organisms study; and part 3 - the
124 potential Interferents applicable to Department of Defense applications.³

125

126 **Part 1:**

127

128 **Environmental Matrix Samples - Aerosol Environmental Matrices**

129

130

131 Method developers shall obtain environmental matrix samples that are representative and
132 consistent with the collection method that is anticipated to ultimately be used in the field. This
133 includes considerations that may be encountered when the collection system is deployed
134 operationally such as collection medium, duration of collection, diversity of geographical areas
135 that will be sampled, climatic/environmental conditions that may be encountered and seasonal
136 changes in the regions of deployment.

137

138 Justifications for the selected conditions that were used to generate the environmental matrix
139 and limitations of the validation based on those criteria must be documented.

140

- 141 • Method developers shall test the environmental matrix samples for interference using
142 samples inoculated with a target biological threat agent sufficient to achieve 95%
143 probability of detection.
- 144 • Cross-reactivity testing will include sufficient samples and replicates to ensure each
145 environmental condition is adequately represented.

146

147

³ Added in June 2015 for the Department of Defense project.

148

149 **Part 2: Environmental Panel Organisms** - This list is comprised of identified organisms from the
150 environment.

151

152 Inclusion of all environmental panel organisms is not a requirement if a method developer provides
153 appropriate justification that the intended use of the assay permits the exclusion of specific panel
154 organisms. Justification for exclusion of any environmental panel organism(s) must be documented
155 and submitted.

156

157 Organisms and cell lines may be tested as isolated DNA, or as pools of isolated DNA. Isolated DNA
158 may be combined into pools of up to 10 panel organisms, with each panel organism represented at
159 10 times the AMDL, where possible. The combined DNA pools are tested in the presence (at 2 times
160 the AMDL) and absence of the target gene or gene fragment. If an unexpected result occurs, each of
161 the individual environmental organisms from a failed pool must be individually re-tested at 10 times
162 the AMDL with and without the target gene or gene fragment at 2x the AMDL in the candidate
163 method DNA elution buffer.

164

165 DNA in this list that already appear in the inclusivity or exclusivity panel do not need to be tested
166 again as part of the environmental factors panel.

167

168 • **Potential bacterial biothreat agents**

169

Bacillus anthracis Ames

170

Yersinia pestis Colorado-92

171

Francisella tularensis subsp. *tularensis* Schu-S4

172

Burkholderia pseudomallei

173

Burkholderia mallei

174

Brucella melitensis

175

176 • **Cultivable bacteria identified as being present in air soil or water**

177

Acinetobacter lwoffii

178

Agrobacterium tumefaciens

179

Bacillus amyloliquefaciens

180

Bacillus cohnii

181

Bacillus psychrosaccharolyticus

182

Bacillus benzoovorans

183

Bacillus megaterium

184

Bacillus horikoshii

185

Bacillus macroides

186

Bacteroides fragilis

187

Burkholderia cepacia

188

Burkholderia gladioli

189

Burkholderia stabilis

190

Burkholderia plantarii

191

Chryseobacterium indologenes

192

Clostridium sardiniense

193

Clostridium perfringens

194

Deinococcus radiodurans

195

Delftia acidovorans

196

Escherichia coli K12

197	<i>Fusobacterium nucleatum</i>
198	<i>Lactobacillus plantarum</i>
199	<i>Legionella pneumophila</i>
200	<i>Listeria monocytogenes</i>
201	<i>Moraxella nonliquefaciens</i>
202	<i>Mycobacterium smegmatis</i>
203	<i>Neisseria lactamica</i>
204	<i>Pseudomonas aeruginosa</i>
205	<i>Rhodobacter sphaeroides</i>
206	<i>Riemerella anatipestifer</i>
207	<i>Shewanella oneidensis</i>
208	<i>Staphylococcus aureus</i>
209	<i>Stenotrophomonas maltophilia</i>
210	<i>Streptococcus pneumoniae</i>
211	<i>Streptomyces coelicolor</i>
212	<i>Synechocystis</i>
213	<i>Vibrio cholerae</i>
214	
215	• Microbial eukaryotes
216	
217	<u>Freshwater amoebae</u>
218	<i>Acanthamoeba castellanii</i>
219	<i>Naegleria fowleri</i>
220	
221	<u>Fungi</u>
222	<i>Alternaria alternata</i>
223	<i>Aspergillus fumigatus</i>
224	<i>Aureobasidium pullulans</i>
225	<i>Cladosporium cladosporioides</i>
226	<i>Cladosporium sphaerospermum</i>
227	<i>Epicoccum nigrum</i>
228	<i>Eurotium amstelodami</i>
229	<i>Mucor racemosus</i>
230	<i>Paecilomyces variotii</i>
231	<i>Penicillium chrysogenum</i>
232	<i>Wallemia sebi</i>
233	
234	

235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279

- **DNA from higher eukaryotes**

- Plant Pollen⁴

- Zea mays* (corn)

- Pinus* spp . (pine)

- Gossypium* spp. (Cotton)

- Arthropods

- Aedes aegypti* (ATCC /CCL-125(tm) mosquito cell line)

- Aedes albopictus* (Mosquito C6/36 cell line)

- Dermatophagoides pteronyssinus* (Dust mite -commercial source)

- Xenopsylla cheopis* Flea (Rocky Mountain labs)

- Drosophila* cell line

- Musca domestica* (housefly) ARS, USDA, Fargo, ND

- Gypsy moth cell lines LED652Y cell line (baculovirus)– Invitrogen

- Cockroach (commercial source)

- Tick (*Amblyomma* and *Dermacentor* tick species for *F. tularensis* detection assays)⁵

- Vertebrates

- Mus musculus* (ATCC/HB-123) mouse

- Rattus norvegicus* (ATCC/CRL-1896) rat

- Canis familiaris*(ATCC/CCL-183) dog

- Felis catus* (ATCC/CRL-8727) cat

- Homo sapiens* (HeLa cell line ATCC/CCL-2) human

- Gallus gallus domesticus* (Chicken)

- Capra hircus* (Goat)⁶

- **Biological insecticides** – Strains of *B. thuringiensis* present in commercially available insecticides have been extensively used in hoaxes and are likely to be harvested in air collectors. For these reasons, it should be used to assess the specificity of these threat assays.

- B. thuringiensis* subsp. *israelensis*

- B. thuringiensis* subsp. *kurstaki*

- B. thuringiensis* subsp. *morrisoni*

- Serenade (Fungicide) *B. subtilis* (QST713)

Viral agents have also been used for insect control. Two representative products are:

- Gypcheck for gypsy moths (*Lymanteria dispar* nuclear polyhedrosis virus)

- Cyd-X for codling moths (Codling moth granulosis virus)

⁴ If pollen is unavailable, vegetative DNA is acceptable

⁵ Added by SPADA on (future approval date).

⁶ Added by SPADA on September 1, 2015

280 **Part 3: Potential Interferents Study**

281

282 The Potential Interferents Study supplements the Environmental Factors Study, and is applicable
283 to all biological threat agent detection assays for Department of Defense applications. Table VI
284 provides a list of potential Interferents that are likely to be encountered in various Department
285 of Defense applications.

286

287 Method developers and evaluators shall determine the most appropriate potential Interferents
288 for their application. Interferents shall be spiked at a final test concentration of 1 µg/ml directly
289 into the sample collection buffer. Sample collection buffers spiked with potential Interferents
290 shall be inoculated at 2 times the AMDL (or AMIL) with one of the target biological threat
291 agents.

292

293 Spiked / inoculated sample collection buffers shall be tested using the procedure specified by
294 the candidate method. A candidate method that fails at the 1 microgram per ml level may be
295 reevaluated at lower concentrations until the inhibition level is determined.

296

297 It is expected that all samples are correctly identified as positive.

298

299

300
301

Table VI: Potential Interferents

Compounds		Potential Theaters of Operation
group 1: petroleum-based	JP-8 ¹	Airfield
	JP-5 ²	Naval
	diesel/gasoline mixture	Ground
	fog oil (standard grade fuel number 2)	naval, ground
	burning rubber ³	ground, airfield
group 2: exhaust	gasoline exhaust	Ground
	jet exhaust	naval, airfield
	diesel exhaust	Ground
group 3: obscurants	terephthalic acid ⁴	Ground
	zinc chloride smoke ⁵	Ground
	solvent yellow 33 ⁶	Ground
group 4: environmental	burning vegetation	ground, airfield
	road dust	Ground
	sea water (sea spray)	Naval
group 5: chemicals	brake fluid ⁷	All
	brake dust ⁸	Ground
	cleaning solvent, MIL-L-63460 ⁹	All
	explosive residues a) high explosives ¹⁰ b) artillery propellant ¹¹	All

302
303
304
305
306
307
308


Table VI is offered for guidance and there are no mandatory minimum requirements for the number of potential Interferents to be tested.

¹ **JP-8.** Air Force formulation jet fuel.

² **JP-5.** A yellow kerosene-based jet fuel with a lower flash point developed for use in aircraft stationed aboard aircraft carriers, where the risk from fire is particularly great. JP-5 is a complex mixture of hydrocarbons, containing alkanes, naphthenes, and aromatic hydrocarbons.


³ **Burning rubber** (tire smoke). Gaseous C1-C5 hydrocarbons: methane; ethane; isopropene; butadiene; propane. Polycyclic aromatic hydrocarbons (58-6800 ng/m³): parabenzo(a)pyrene; polychlorinated dibenzo-p-dioxins (PCDD); polychlorinated dibenzofurans (PCDF). Metals (0.7 - 8 mg/m³): zinc; lead; cadmium.

⁴ **Terephthalic acid.** Used in the AN/M83 hand grenade currently used by US military.



⁵ **Zinc chloride smoke.** Also known as “zinc chloride smoke” and “HC smoke”. Was used in the M8 grenade and still used in 155mm artillery shells. HC smoke is composed of 45% hexachloroethane, 45% zinc oxide, and 10% aluminum.

⁶ **Solvent yellow 33** [IUPAC name: 2-(2-quinolyyl)-1,3-indandione] is a new formulation being developed for the M18 grenade.



⁷ **Brake fluid.** DOT 4 is primarily composed of glycol and borate esters. DOT 5 is silicone-based brake fluid. The main difference is that DOT 4 is hydroscopic whereas DOT 5 is hydrophobic. DOT 5 is often used in military vehicles because it is more stable over time requires less maintenance

⁸ **Brake dust.** Fe particles caused by abrasion of the cast iron brake rotor by the pad and secondly fibers from the semi metallic elements of the brake pad. The remainder of the dust residue is carbon content within the brake pad.

⁹ **MIL-L-63460,** "Military Specification, Lubricant, Cleaner and Preservative for Weapons and Weapons Systems"; trade name "Break-Free CLP". Hyperlink: [Midway USA](#).

¹⁰ **High explosives.** The M795 155mm projectile is the US Army / Marine Corp's current standard projectile containing 10.8 kg of TNT. The M795 projectile replaced the M107 projectile that contained Composition B which is a 60/40 mixture of RDX/TNT. RDX is cyclotrimethylene trinitramine. Suggestion: test RDX/TNT together.

¹¹ **Artillery propellant.** Modern gun propellants are divided into three classes: single-base propellants which are mainly or entirely nitrocellulose based, double-base propellants composed of a combination of nitrocellulose and nitroglycerin, and triple base composed of a combination of nitrocellulose and nitroglycerin and nitroguanidine. Suggestion: test total nitrocellulose/ nitroglycerin nitroguanidine together.



AOAC Stakeholder Panel on Agent Detection Assays

Yersinia pestis Working Group



Luther Lindler
Chief Scientist (ST)
Chemical and Biological Defense Division
Homeland Security Advanced Research
Projects Agency
Science and Technology Directorate

Fitness for Purpose

“Field-deployable detection of *Yersinia pestis* in samples from Department of Defense aerosol collection devices.”



SPADA *Yersinia pestis* Working Group Members

Luther Lindler, DHS (Chair)
Jessica Appler, HHS/BARDA
Jeff Ballin, ECBC
Timothy Bauer, NSWC
Ryan Cahall, Censeo Insight
Ken Damer, NGC
Matthew Davenport, DHS
Christina Egan, NYSDOH
Jennifer Gibbons, ECBC
Ted Hadfield, Hadeco
Malcolm Johns, DHS
Matthew Lesho, Luminex

Laura Maple, NSWC
Traci Pals, DTRA
Robert Perry, University of Kentucky
Michael Retford, JBTS JPM NBCCA
Michael Scheckelhoff, DHS
Sanjiv Shah, EPA
Shanmuga Sozhamannan, CRP
David Wagner, NAU
David Watson, DTRA
Patricia Worsham, USAMRIID



Yersinia pestis Working Group *Work to Date*

- Working Group Launch (September, 2015)
- 3 teleconferences (October 2015 – November 2015)
- 1 SMPR Drafted
- Public comment period (January 8, 2016 – February 5, 2016)
- SMPR made ready for SPADA review and approval




Background

- *Yersinia pestis* and *Yersinia pseudotuberculosis* are closest related
- *Y. pestis* generally has 2 species specific plasmids (“pestoides F” exception)
- Highly monomorphic but does show gene loss/gain, intra-genome recombination events, SNPs
- Genetic relatedness around phenotypic groups (PE, ANT, MED, ORI)
- Goal- to choose available examples from current clades



Genetic Relatedness



Morelli et al. 2010. Nature Genetics

Method Performance Requirements

Parameter	Minimum Performance Requirement
AMDL	2,000 standardized cells of <i>Yersinia pestis</i> strain CO92 per mL liquid in the candidate method sample collection buffer.
Probability of Detection at AMDL within sample collection buffer	≥ 0.95
Probability of Detection at AMDL in environmental matrix materials.	≥ 0.95
System False-Negative Rate using spiked environmental matrix materials.	≤ 5%
System False-Positive Rate using environmental matrix materials.	≤ 5%
Inclusivity	All inclusivity strains (Table III) must test positive at 2x the AMDL †
Exclusivity	All exclusivity strains (Table IV and Table V; part 2) must test negative at 10x the AMDL †
Notes: † 100% correct analyses are expected. All discrepancies are to be retested following the AOAC Guidelines for Validation of Biological Threat Agent Methods and/or Procedures. 1	

SMPR Key Points Inclusivity Panel

Number	Strain	Achtman Genotype	Comments	Availability
1	CO92	1.ORI.c	Well studied example of epidemic strain of pestis, recent isolate	CDC, WRAIR, RIID
2	KIM	2.Med	Well studied strain in academic circles, virulence data extensive	CDC, WRAIR, RIID
3	Antiqua	1.Ant b	Ancient strain near root of tree	CDC, WRAIR, RIID
4	Pestoides B	0.PE1		CDC, WRAIR, RIID
5	Pestoides F	0.PE2.a	pPst negative, old strain in terms of phylogeny	CDC, WRAIR, RIID
6	Pestoides G	0.PE2.b	pPst negative	CDC, WRAIR, RIID
7	Angola	0.PE3	A "pestoides" in everything except name	CDC, WRAIR, RIID
8	Nairobi	1.Ant a		CDC, WRAIR, RIID
9	Harbin35	2 Ant	Rumored to be used or resulted from infection during experiments by Japanese BW Unit 731	CDC, WRAIR, RIID
10	PBM19	1.ORI.a		CDC, WRAIR, RIID
11	Java9	1.ORI	pFra negative	CDC, WRAIR, RIID
12	A1122	1.ORI.a	Well characterized US isolate that is pgm- and pCD-; also has 2X large pPst plasmid	CDC, WRAIR, RIID
13	Nicholsk 41	2.ANT		CDC, WRAIR, RIID
14	Shasta	1.ORI	YE0387; Shasta (20 Oct 54); Shasta; human case; USA: Ca; 1960 6LY; UCC YERS074	CDC, RIID
15	Dodson	1.ORI	Dodson (Aug 70); human case: male age 4.5 years; USA: Arizona (Tuba City); 27 Jun 67; UCC YERS073	CDC, RIID
16	El Dorado	1.Ori		CDC

SMPR Key Points

Exclusivity Panel

	Species	Strain		Comments	Availab
YPNN1	<i>Yersinia ruckeri</i>	YERS063			RIID
YPNN2	<i>Yersinia rohdei</i>	YERS062			RIID
YPNN3	<i>Yersinia pseudotuberculosis</i>	PB1/+	1	sequenced	WRAIR
YPNN4	<i>Yersinia pseudotuberculosis</i>	IP32953	1	sequenced	WRAIR
YPNN5	<i>Yersinia pseudotuberculosis</i>	YPIII	3	sequenced	WRAIR
YPNN6	<i>Yersinia pseudotuberculosis</i>	Pa3606	1b		WRAIR
YPNN7	<i>Yersinia pseudotuberculosis</i>	IB	1b		WRAIR
YPNN8	<i>Yersinia pseudotuberculosis</i>	EP2/+	1		WRAIR
YPNN9	<i>Yersinia pseudotuberculosis</i>	MD67	1		WRAIR
YPNN10	<i>Yersinia pseudotuberculosis</i>	1	1a		WRAIR
YPNN11	<i>Yersinia enterocolitica</i>	WA	O:8		WRAIR
YPNN12	<i>Yersinia enterocolitica</i>	8081	O:8	sequenced	WRAIR
YPNN13	<i>Yersinia enterocolitica</i>	2516-87	O:9		WRAIR
YPNN14	<i>Yersinia kirstensenii</i>	Y231		non-pathogenic	WRAIR
YPNN15	<i>Yersinia frederiksenii</i>	Y225		non-pathogenic	WRAIR
YPNN16	<i>Yersinia intermedia</i>	Y228		non-pathogenic	WRAIR
YPNN17	<i>Yersinia aldovae</i>	670-83		non-pathogenic	WRAIR

SMPR Key Points

- Environmental Panel same as SPADA 1
- Interferents include military relevant chemicals (jet fuel, Terephthalic acid found in hand grenades, cleaning solvent, MIL-L-63460, dust, diesel exhaust, artillery propellant, etc.)



Comments Submitted (if any)

- No comments submitted for this SMPR



Motion

- Motion to accept the Standard Method Performance Requirements for *Yersinia pestis* as presented.



Discussion?



Homeland Security

Science and Technology

3
4 **Standard Method Performance Requirements (SMPRs®) for**
5 **DNA-based methods of detecting *Yersinia pestis* in field-deployable, Department of Defense**
6 **aerosol collection devices**

7
8 **Intended Use:** *Field-deployed use for analysis of aerosol collection filters and/or liquids*

9
10 **1. Applicability:** Detection of *Yersinia pestis* in collection buffers from aerosol collection
11 devices. Field-deployable assays are preferred.

12
13 **2. Analytical Technique:** Molecular detection of nucleic acid.

14
15 **3. Definitions:**

16
17 **Acceptable Minimum Detection Level (AMDL)**

18 The predetermined minimum level of an analyte, as specified by an expert committee which
19 must be detected by the candidate method at a specified probability of detection (POD).

20
21 **Exclusivity**

22 Study involving pure non-target strains, which are potentially cross-reactive, that shall not
23 be detected or enumerated by the candidate method.

24
25 **Inclusivity**

26 Study involving pure target strains that shall be detected or enumerated by the candidate
27 method.

28
29 **Maximum Time-To- Result**

30 Maximum time to complete an analysis starting from the collection buffer ~~test portion~~
31 ~~preparation~~ to assay result.

32
33 **Probability of Detection (POD)**

34 The proportion of positive analytical outcomes for a qualitative method for a given matrix at
35 a specified analyte level or concentration with a ≥ 0.95 confidence interval.

36
37 **System False Negative Rate**

38 Proportion of test results that are negative contained within a population of known
39 positives

40
41 **System False Positive Rate**

42 Proportion of test results that are positive contained within a population of known
43 negatives.

44
45
46 **4. Method Performance Requirements:**

47
48 See Table I.

50 **5. System suitability tests and/or analytical quality control:**

51 The controls listed in Table II shall be embedded in assays as appropriate. Manufacturer
52 must provide written justification if controls are not embedded in the assay.

53
54 **6. Validation Guidance:**

55 AOAC INTERNATIONAL Methods Committee Guidelines for Validation of Biological Threat
56 Agent Methods and/or Procedures (AOAC INTERNATIONAL Official Methods of Analysis,
57 2012, Appendix I).

58
59 Inclusivity and exclusivity panel organisms used for evaluation must be characterized and
60 documented to truly be the species and strains they are purported to be.

61
62 **8. Maximum time-to-results:** Within four hours.

63
64
65 **Table I: Method Performance Requirements**

66

Parameter	Minimum Performance Requirement
AMDL	2,000 standardized cells of <i>Yersinia pestis</i> strain CO92 per mL liquid in the candidate method sample collection buffer.
Probability of Detection at AMDL within sample collection buffer	≥ 0.95
Probability of Detection at AMDL in environmental matrix materials.	≥ 0.95
System False-Negative Rate using spiked environmental matrix materials.	≤ 5%
System False-Positive Rate using environmental matrix materials.	≤ 5%
Inclusivity	All inclusivity strains (Table III) must test positive at 2x the AMDL [†]
Exclusivity	All exclusivity strains (Table IV and Table V; part 2) must test negative at 10x the AMDL [†]
Notes: † 100% correct analyses are expected. All discrepancies are to be retested following the AOAC Guidelines for Validation of Biological Threat Agent Methods and/or Procedures. ¹	

67
68
¹ Official Methods of Analysis of AOAC INTERNATIONAL (2012) 19th Ed., AOAC INTERNATIONAL, Gaithersburg, MD, USA, APPENDIX I; also on-line at http://www.eoma.aoc.org/app_i.pdf.

TABLE II: Controls

Control	Description	Implementation
Positive Control	<p>This control is designed to demonstrate an appropriate test response. The positive control should be included at a low but easily detectable concentration, and should monitor the performance of the entire assay. The purpose of using a low concentration of positive control is to demonstrate that the assay sensitivity is performing at a previously determined level of sensitivity.</p>	<p>Single use per sample (or sample set) run</p>
Negative Control	<p>This control is designed to demonstrate that the assay itself does not produce a detection in the absence of the target organism. The purpose of this control is to rule-out causes of false positives, such as contamination in the assay or test.</p>	<p>Single use per sample (or sample set) run</p>
Inhibition Control	<p>This control is designed to specifically address the impact of a sample or sample matrix on the assay's ability to detect the target organism.</p>	<p>Single use per sample (or sample set) run</p>

71
72
73

Table III: Inclusivity Panel

Number	Strain	Achtman Genotype	Comments	Availability
1	CO92	1.ORI.c	Well studied example of epidemic strain of pestis, recent isolate	CDC, WRAIR, RIID
2	KIM	2.Med	Well studied strain in academic circles, virulence data extensive	CDC, WRAIR, RIID
3	Antiqua	1.Ant b	Ancient strain near root of tree	CDC, WRAIR, RIID
4	Pestoides B	0.PE1		CDC, WRAIR, RIID
5	Pestoides F	0.PE2.a	pPst negative, old strain in terms of phylogeny	CDC, WRAIR, RIID
6	Pestoides G	0.PE2.b	pPst negative	CDC, WRAIR, RIID
7	Angola	0.PE3	A "pestoides" in everything except name	CDC, WRAIR, RIID
8	Nairobi	1.Ant a		CDC, WRAIR, RIID
9	Harbin35	2 Ant	Rumored to be used or resulted from infection during experiments by Japanese BW Unit 731	CDC, WRAIR, RIID
10	PBM19	1.ORI.a		CDC, WRAIR, RIID
11	Java9	1.ORI	pFra negative	CDC, WRAIR, RIID
12	A1122	1.ORI.a	Well characterized US isolate that is pgm- and pCD-; also has 2X large pPst plasmid	CDC, WRAIR, RIID
13	Nicholisk 41	2.ANT		CDC, WRAIR, RIID
14	Shasta	1.ORI	YE0387; Shasta (20 Oct 54); Shasta; human case; USA: Ca; 1960 6LY; UCC YERS074	CDC, RIID
15	Dodson	1.ORI	Dodson (Aug 70); human case: male age 4.5 years; USA: Arizona (Tuba City); 27 Jun 67; UCC YERS073	CDC, RIID
16	El Dorado			

74
75

Note on plasmid nomenclature: pMT1 = pFRA; pPCP1 = pPST = pPLA; pCD1 = pYB = pCAD.

76
77
78
79

Table IV: Exclusivity Panel (near-neighbor)

	Species	Strain		Comments	Availability
YPNN1	<i>Yersinia ruckeri</i>	YERS063			RIID
YPNN2	<i>Yersinia rohdei</i>	YERS062			RIID
YPNN3	<i>Yersinia pseudotuberculosis</i>	PB1/+	1	sequenced	WRAIR
YPNN4	<i>Yersinia pseudotuberculosis</i>	IP32953	1	sequenced	WRAIR
YPNN5	<i>Yersinia pseudotuberculosis</i>	YPIII	3	sequenced	WRAIR
YPNN6	<i>Yersinia pseudotuberculosis</i>	Pa3606	1b		WRAIR
YPNN7	<i>Yersinia pseudotuberculosis</i>	IB	1b		WRAIR
YPNN8	<i>Yersinia pseudotuberculosis</i>	EP2/+	1		WRAIR
YPNN9	<i>Yersinia pseudotuberculosis</i>	MD67	1		WRAIR
YPNN10	<i>Yersinia pseudotuberculosis</i>	1	1a		WRAIR
YPNN11	<i>Yersinia enterocolitica</i>	WA	O:8		WRAIR
YPNN12	<i>Yersinia enterocolitica</i>	8081	O:8	sequenced	WRAIR
YPNN13	<i>Yersinia enterocolitica</i>	2516-87	O:9		WRAIR
YPNN14	<i>Yersinia kirstensenii</i>	Y231		non-pathogenic	WRAIR
YPNN15	<i>Yersinia frederiksenii</i>	Y225		non-pathogenic	WRAIR
YPNN16	<i>Yersinia intermedia</i>	Y228		non-pathogenic	WRAIR
YPNN17	<i>Yersinia aldovae</i>	670-83		non-pathogenic	WRAIR

80
81
82
83
84
85
86
87
88
89

Guidance

Organisms may be tested as isolated DNA, or combined to form a pool of isolated DNA. Isolated DNA may be combined into pools of up to 10 exclusivity panel organisms, with each panel organism represented at 10 times the AMDL, where possible. If an unexpected result occurs, each of the exclusivity organisms from a failed pool must be individually re-tested at 10 times the AMDL.

90 **Table V: Environmental Factors For Validating Biological Threat Agent Detection Assays**

91
92 [Adapted from the Environmental Factors Panel approved by SPADA on June 10, 2010.]

93
94 The Environmental Factors Studies supplement the biological threat agent near-neighbor
95 exclusivity testing panel. There are three parts to Environmental Factors studies: part 1 -
96 environmental matrix samples; part 2 - the environmental organisms study; and part 3 - the
97 potential interferants applicable to Department of Defense applications.² **Part 2 is not**
98 **applicable to techniques that do not detect nucleic acid.**
99

100
101 **Part 1:**

102
103 **Environmental Matrix Samples - Aerosol Environmental Matrices**

104
105
106 Method developers shall obtain environmental matrix samples that are representative and
107 consistent with the collection method that is anticipated to ultimately be used in the field. This
108 includes considerations that may be encountered when the collection system is deployed
109 operationally such as collection medium, duration of collection, diversity of geographical areas
110 that will be sampled, climatic/environmental conditions that may be encountered and seasonal
111 changes in the regions of deployment.

112
113 Justifications for the selected conditions that were used to generate the environmental matrix
114 and limitations of the validation based on those criteria must be documented.

- 115
116 • Method developers shall test the environmental matrix samples for interference using
117 samples inoculated with a target biological threat agent sufficient to achieve 95%
118 probability of detection.
119 • Cross-reactivity testing will include sufficient samples and replicates to ensure each
120 environmental condition is adequately represented.
121
122

² Added in June 2015 for the Department of Defense project.

123

124 **Part 2: Environmental Panel Organisms** - This list is comprised of identified organisms from the
125 environment.

126

127 Inclusion of all environmental panel organisms is not a requirement if a method developer
128 provides appropriate justification that the intended use of the assay permits the exclusion of
129 specific panel organisms. Justification for exclusion of any environmental panel organism(s)
130 must be documented and submitted.

131

132 Organisms and cell lines may be tested as isolated DNA, or as pools of isolated DNA. Isolated
133 DNA may be combined into pools of up to 10 panel organisms, with each panel organism
134 represented at 10 times the AMDL, where possible. The combined DNA pools are tested in the
135 presence (at 2 times the AMDL) and absence of the target gene or gene fragment. If an
136 unexpected result occurs, each of the individual environmental organisms from a failed pool
137 must be individually re-tested at 10 times the AMDL with and without the target gene or gene
138 fragment at 2x the AMDL in the candidate method DNA elution buffer.

139

140 DNA in this list that already appear in the inclusivity or exclusivity panel do not need to be
141 tested again as part of the environmental factors panel.

142

143 • **Potential bacterial biothreat agents**

144 *Bacillus anthracis* Ames

145 *Yersinia pestis* Colorado-92

146 *Francisella tularensis* subsp. *tularensis* Schu-S4

147 *Burkholderia pseudomallei*

148 *Burkholderia mallei*

149 *Brucella melitensis*

150

151 • **Cultivable bacteria identified as being present in air soil or water**

152 *Acinetobacter lwoffii*

153 *Agrobacterium tumefaciens*

154 *Bacillus amyloliquefaciens*

155 *Bacillus cohnii*

156 *Bacillus psychrosaccharolyticus*

157 *Bacillus benzoovorans*

158 *Bacillus megaterium*

159 *Bacillus horikoshii*

160 *Bacillus macroides*

161 *Bacteroides fragilis*

162 *Burkholderia cepacia*

163 *Burkholderia gladioli*

164 *Burkholderia stabilis*

165 *Burkholderia plantarii*

166 *Chryseobacterium indologenes*

167 *Clostridium sardiniense*

168 *Clostridium perfringens*

169 *Deinococcus radiodurans*

170 *Delftia acidovorans*

171 *Escherichia coli* K12

172	<i>Fusobacterium nucleatum</i>
173	<i>Lactobacillus plantarum</i>
174	<i>Legionella pneumophila</i>
175	<i>Listeria monocytogenes</i>
176	<i>Moraxella nonliquefaciens</i>
177	<i>Mycobacterium smegmatis</i>
178	<i>Neisseria lactamica</i>
179	<i>Pseudomonas aeruginosa</i>
180	<i>Rhodobacter sphaeroides</i>
181	<i>Riemerella anatipestifer</i>
182	<i>Shewanella oneidensis</i>
183	<i>Staphylococcus aureus</i>
184	<i>Stenotrophomonas maltophilia</i>
185	<i>Streptococcus pneumoniae</i>
186	<i>Streptomyces coelicolor</i>
187	<i>Synechocystis</i>
188	<i>Vibrio cholerae</i>
189	
190	• Microbial eukaryotes
191	
192	<u>Freshwater amoebae</u>
193	<i>Acanthamoeba castellanii</i>
194	<i>Naegleria fowleri</i>
195	
196	<u>Fungi</u>
197	<i>Alternaria alternata</i>
198	<i>Aspergillus fumigatus</i>
199	<i>Aureobasidium pullulans</i>
200	<i>Cladosporium cladosporioides</i>
201	<i>Cladosporium sphaerospermum</i>
202	<i>Epicoccum nigrum</i>
203	<i>Eurotium amstelodami</i>
204	<i>Mucor racemosus</i>
205	<i>Paecilomyces variotii</i>
206	<i>Penicillium chrysogenum</i>
207	<i>Wallemia sebi</i>
208	
209	

210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254

- **DNA from higher eukaryotes**

- Plant Pollen³

- Zea mays* (corn)

- Pinus* spp . (pine)

- Gossypium* spp. (Cotton)

- Arthropods

- Aedes aegypti* (ATCC /CCL-125(tm) mosquito cell line)

- Aedes albopictus* (Mosquito C6/36 cell line)

- Dermatophagoides pteronyssinus* (Dust mite -commercial source)

- Xenopsylla cheopis* Flea (Rocky Mountain labs)

- Drosophila* cell line

- Musca domestica* (housefly) ARS, USDA, Fargo, ND

- Gypsy moth cell lines LED652Y cell line (baculovirus)– Invitrogen

- Cockroach (commercial source)

- Tick (*Amblyomma* and *Dermacentor* tick species for *F. tularensis* detection assays)⁴

- Vertebrates

- Mus musculus* (ATCC/HB-123) mouse

- Rattus norvegicus* (ATCC/CRL-1896) rat

- Canis familiaris*(ATCC/CCL-183) dog

- Felis catus* (ATCC/CRL-8727) cat

- Homo sapiens* (HeLa cell line ATCC/CCL-2) human

- Gallus gallus domesticus* (Chicken)

- Goat⁵

- **Biological insecticides** – Strains of *B. thuringiensis* present in commercially available insecticides have been extensively used in hoaxes and are likely to be harvested in air collectors. For these reasons, it should be used to assess the specificity of these threat assays.

- B. thuringiensis* subsp. *israelensis*

- B. thuringiensis* subsp. *kurstaki*

- B. thuringiensis* subsp. *morrisoni*

- Serenade (Fungicide) *B. subtilis* (QST713)

Viral agents have also been used for insect control. Two representative products are:

Gypcheck for gypsy moths (*Lymanteria dispar* nuclear polyhedrosis virus)

Cyd-X for codling moths (Codling moth granulosis virus)

³ If pollen is unavailable, vegetative DNA is acceptable

⁴ Added by SPADA on (future approval date).

⁵ Added by SPADA on September 1, 2015

255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279

Part 3: Potential Interferants Study

The Potential Interferants Study supplements the Environmental Factors Study, and is applicable to all biological threat agent detection assays for Department of Defense applications. Table VI provides a list of potential interferants that are likely to be encountered in various Department of Defense applications.

Method developers and evaluators shall determine the most appropriate potential interferants for their application. Interferants shall be spiked at a final test concentration of 1 µg/ml directly into the sample collection buffer. Interferants may be pooled. Sample collection buffers spiked with potential interferants shall be inoculated at 2 times the AMDL (or AMIL) with one of the target biological threat agents.

Spiked / inoculated sample collection buffers shall be tested using the procedure specified by the candidate method. A candidate method that fails at the 1 microgram per ml level may be reevaluated at lower concentrations until the inhibition level is determined.

It is expected that all samples are correctly identified as positive. If using pooled samples of potential interferants, and a negative result occurs, then the pooled potential interferants shall be tested separately at the 2 times the AMDL (or AMIL) with one of the target biological threat agents.

280
281

Table VI: Potential Interferants

Compounds		Potential Theaters of Operation
group 1: petroleum-based	JP-8 ¹	airfield
	JP-5 ²	naval
	diesel/gasoline mixture	ground
	fog oil (standard grade fuel number 2)	naval, ground
	burning rubber ³	ground, airfield
group 2: exhaust	gasoline exhaust	ground
	jet exhaust	naval, airfield
	diesel exhaust	ground
group 3: obscurants	terephthalic acid ⁴	ground
	zinc chloride smoke ⁵	ground
	solvent yellow 33 ⁶	ground
group 4: environmental	burning vegetation	ground, airfield
	road dust	ground
	sea water (sea spray)	naval
group 5: chemicals	brake fluid ⁷	all
	brake dust ⁸	ground
	cleaning solvent, MIL-L-63460 ⁹	all
	explosive residues a) high explosives ¹⁰ b) artillery propellant ¹¹	all

282
283
284

285
286
287
288


Table VI is offered for guidance and there are no mandatory minimum requirements for the number of potential interferants to be tested.

¹ **JP-8.** Air Force formulation jet fuel.

² **JP-5.** A yellow kerosene-based jet fuel with a lower flash point developed for use in aircraft stationed aboard aircraft carriers, where the risk from fire is particularly great. JP-5 is a complex mixture of hydrocarbons, containing alkanes, naphthenes, and aromatic hydrocarbons.


³ **Burning rubber** (tire smoke). Gaseous C1-C5 hydrocarbons: methane; ethane; isopropene; butadiene; propane. Polycyclic aromatic hydrocarbons (58-6800 ng/m³): parabenzo(a)pyrene; polychlorinated dibenzo-p-dioxins (PCDD); polychlorinated dibenzofurans (PCDF). Metals (0.7 - 8 mg/m³): zinc; lead; cadmium.

⁴ **Terephthalic acid.** Used in the AN/M83 hand grenade currently used by US military.



⁵ **Zinc chloride smoke.** Also known as “zinc chloride smoke” and “HC smoke”. Was used in the M8 grenade and still used in 155mm artillery shells. HC smoke is composed of 45% hexachloroethane, 45% zinc oxide, and 10% aluminum.

⁶ **Solvent yellow 33** [IUPAC name: 2-(2-quinoly)-1,3-indandione] is a new formulation being developed for the M18 grenade.




⁷ **Brake fluid.** DOT 4 is the most common brake fluid, primarily composed of glycol and borate esters. DOT 5 is silicone-based brake fluid. The main difference is that DOT 4 is hygroscopic whereas DOT 5 is hydrophobic. DOT 5 is often used in military vehicles because it is more stable over time requires less maintenance

⁸ **Brake dust.** Fe particles caused by abrasion of the cast iron brake rotor by the pad and secondly fibers from the semi metallic elements of the brake pad. The remainder of the dust residue is carbon content within the brake pad.

⁹ **MIL-L-63460,** "Military Specification, Lubricant, Cleaner and Preservative for Weapons and Weapons Systems"; trade name “Break-Free CLP”. Hyperlink: [Midway USA](#).

¹⁰ **High explosives.** The M795 155mm projectile is the US Army / Marine Corp's current standard projectile containing 10.8 kg of TNT. The M795 projectile replaced the M107 projectile that contained Composition B which is a 60/40 mixture of RDX/TNT. RDX is cyclotrimethylene trinitramine. Suggestion: test RDX/TNT together.

¹¹ **Artillery propellant.** Modern gun propellants are divided into three classes: single-base propellants which are mainly or entirely nitrocellulose based, double-base propellants composed of a combination of nitrocellulose and nitroglycerin, and triple base composed of a combination of nitrocellulose and nitroglycerin and nitroguanidine. Suggestion: test total nitrocellulose/ nitroglycerin nitroguanidine together.



AOAC INTERNATIONAL STAKEHOLDER PANEL ON AGENT DETECTION ASSAYS

Francisella tularensis Working Group
Co-Chairs: Paul Keim and David Wagner, NAU
SMPR Presentation
March 22, 2016

Rockville, Maryland, USA

Fitness for Purpose

Field-deployable detection of
Francisella tularensis in samples
from Department of Defense aerosol
collection devices.



**SPADA *Francisella tularensis*
Working Group Members**

Paul Keim, NAU	Katalin Kiss, ATCC
David Wagner, NAU	Laura Maple, NSWC
Mats Forsman, FOI	Stephen Morse, CDC
Jessica Appler, HHS BARDA	Pejman Naraghi-Arani,
Timothy Bauer, NSWC	InSilixa
Linda Beck, NSWC	Denise Pettit, State of NC
Julie Boylan, DTRA	Michael Retford, JBTDS JPM
Ryan Cahall, Censeo Insight	NBCCA
Ken Damer, Northrop	Mark Scheckelhoff, DHS
Grumman	Sanjiv Shah, US EPA
Matthew Davenport, DHS	Shanmuga Sozhamannan,
Steven Hinrichs, UNMC	CRP




***Francisella tularensis* Working Group
Work to Date**

- Working Group Launch (September, 2015)
- Four teleconferences (November 2015 – December 2015)
- One SMPR Drafted
- Public comment period (January 8, 2016 – February 5, 2016)
- SMPRs made ready for SPDS review and approval




Three *F. tularensis* subspecies



From Keim et al. 2007


Differences among subspecies

- *F. tularensis* subsp. *tularensis* (Type A): Common with high diversity – found naturally only in North America
- *F. tularensis* subsp. *holarctica* (Type B): Common with low diversity – found naturally throughout the northern hemisphere
- *F. tularensis* subsp. *mediasiatica*: Rare and reported only occasionally from Central Asia



From Keim et al. 2007

Two distinct groups within Type A




- Type A.I. found in eastern North America and west coast
- Type A.II. found in western North America
- These distributions are correlated with the distributions of specific mammal hosts and tick/fly vectors

From Farlow et al. 2005




Three major subgroups in A.I



Tree from Birdsell et al. 2014
 Mouse data from Molins et al. 2010




Distribution of A.I subgroups in US




From Birdsell et al. 2014

Type B phylogeography circa 2009




From Vogler et al. 2009

Significant diversity in China




From Wang et al. 2014

Significant diversity in Turkey




From Kilic et al. 2015


Current known Type B phylogeography



Identical strains across space & time




Stars = outbreak strains
Circles = archival strains



From Johansson et al. 2014




Near neighbors: the diagnostic challenge



To avoid both false positives and false negatives, diagnostic assays need to target genomic features in space **B**




Sequenced near neighbors



Whole genome SNP phylogeny of *Francisella* by the Swedish Defense Research Agency (FOI)




Many near neighbors are not sequenced



SMPR Key Points

- Inclusivity panel contains representatives from all three subspecies
 - *F. tularensis* subsp. *tularensis* (Type A)
 - Includes both A.I and A.II
 - *F. tularensis* subsp. *holarctica* (Type B)
 - *F. tularensis* subsp. *mediasiatica*
- Exclusivity panel contains representatives from just three near neighbor species
 - *F. novicida*
 - *F. philomiragia*
 - *F. hispaniensis*

Sequenced near neighbors



Whole genome SNP phylogeny of *Francisella* by the Swedish Defense Research Agency (FOI)

SMPR Key Points

- *In silico* screening required
 - Purpose: Predict specificity and inclusivity across all sequenced *Francisella* strains
 - Approach: Use *Basic Alignment Search Tool* (BLAST) to predict hybridization events between signature components (e.g., oligo primers/probes/amplicons) and available *Francisella* genomic sequences in GenBank
 - Include results of *in silico* analyses in method/assay performance evaluation reports

Comments Submitted (if any)

- No comments submitted.



Motion

- Motion to accept the Standard Method Performance Requirements for *Francisella tularensis* as presented.



Discussion?



4 **Standard Method Performance Requirements (SMPRs®) for Detection of *Francisella***
5 ***tularensis* in aerosol collection devices**
6

7 **Intended Use:** Laboratory or field use by Department of Defense trained operators
8

9 **1. Applicability:** Detection of *Francisella tularensis* in collection buffers from aerosol
10 collection devices. Field-deployable assays are preferred.
11

12 **2. Analytical Technique:** Molecular detection of nucleic acid.
13

14 **3. Definitions:**
15

16 **Acceptable Minimum Detection Level (AMDL)**

17 The predetermined minimum level of an analyte, as specified by an expert committee which
18 must be detected by the candidate method at a specified probability of detection (POD).
19

20 **Environmental Factors**

21 For the purposes of this SMPR: any factor in the operating environment of an analytical
22 method, whether abiotic or biotic, that might influence the results of the method.
23

24 **Exclusivity**

25 Study involving pure non-target strains, which are potentially cross-reactive, that shall not
26 be detected or enumerated by the candidate method.
27

28 **Inclusivity**

29 Study involving pure target strains that shall be detected or enumerated by the candidate
30 method.
31

32 **Interferents**

33 A . . . substance in analytical procedures . . . that, at the given concentration, causes a
34 systematic error in the analytical result.¹ Sometimes also known as interferants.
35

36 **Maximum Time-To- Result**

37 Maximum time to complete an analysis starting from the collection buffer to assay result.
38

39 **Probability of Detection (POD)**

40 The proportion of positive analytical outcomes for a qualitative method for a given matrix at
41 a specified analyte level or concentration with a ≥ 0.95 confidence interval.
42
43

¹ International Union Of Pure And Applied Chemistry Analytical Chemistry Division Commission On Analytical Reactions And Reagents* Definition And Classification Of Interferences In Analytical Procedures Prepared For Publication By W. E. Van Der Linden. Pure & Appl. Chem., Vol. 61, No. 1, pp. 91-95, 1989. Printed in Great Britain. © 1989 IUPAC

44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76

System False Negative Rate

Proportion of test results that are negative contained within a population of known positives

System False Positive Rate

Proportion of test results that are positive contained within a population of known negatives.

4. Method Performance Requirements:

See Table I.

5. System suitability tests and/or analytical quality control:

The controls listed in Table II shall be embedded in assays as appropriate. Manufacturer must provide written justification if controls are not embedded in the assay.

6. Validation Guidance: AOAC INTERNATIONAL Methods Committee Guidelines for Validation of Biological Threat Agent Methods and/or Procedures (AOAC INTERNATIONAL Official Methods of Analysis, 2012, Appendix I).

Inclusivity and exclusivity panel organisms used for evaluation must be characterized and documented to truly be the species and strains they are purported to be.

8. Maximum time-to-results: Within four hours.

77
78

Table I: Method Performance Requirements

Parameter	Minimum Performance Requirement
AMDL	2,000 standardized cells per mL liquid in the candidate method sample collection buffer.
Probability of Detection at AMDL within sample collection buffer	≥ 0.95
Probability of Detection at AMDL in environmental matrix materials.	≥ 0.95
System False-Negative Rate using spiked environmental matrix materials.	$\leq 5\%$
System False-Positive Rate using environmental matrix materials.	$\leq 5\%$
Inclusivity	All inclusivity strains (Table III) must test positive at 2x the AMDL [†]
Exclusivity	All exclusivity strains (Table IV and Annex 1 - part 2) must test negative at 10x the AMDL [†]
Notes: [†] 100% correct analyses are expected. All discrepancies are to be re-tested following the AOAC Guidelines for Validation of Biological Threat Agent Methods and/or Procedures ² .	

79
80

² Official Methods of Analysis of AOAC INTERNATIONAL (2012) 19th Ed., AOAC INTERNATIONAL, Gaithersburg, MD, USA, APPENDIX I; also on-line at http://www.eoma.aoac.org/app_i.pdf.

Control	Description	Implementation
Positive Control	This control is designed to demonstrate an appropriate test response. The positive control should be included at a low but easily detectable concentration, and should monitor the performance of the entire assay. The purpose of using a low concentration of positive control is to demonstrate that the assay sensitivity is performing at a previously determined level of sensitivity.	Single use per sample (or sample set) run
Negative Control	This control is designed to demonstrate that the assay itself does not produce a detection in the absence of the target organism. The purpose of this control is to rule-out causes of false positives, such as contamination in the assay or test.	Single use per sample (or sample set) run
Inhibition Control	This control is designed to specifically address the impact of a sample or sample matrix on the assay's ability to detect the target organism.	Single use per sample (or sample set) run

83
84
85
86

Table III: Inclusivity Panel

No.	UCC ^a ID	Genus and species	Strain	Characteristics
1	FRAN001	<i>Francisella tularensis</i>	subsp. <i>tularensis</i>	Type A2 (Type strain)
2	FRAN004	<i>Francisella tularensis</i>	subsp. <i>holarctica</i> (LVS)	Type B (Russian)
3	FRAN012	<i>Francisella tularensis</i>	subsp. <i>holarctica</i>	Type B (United States)
4	FRAN016	<i>Francisella tularensis</i>	subsp. <i>tularensis</i> (SCHU S4)	Type A1 (United States)
5	FRAN024	<i>Francisella tularemia</i>	subsp. <i>holarctica</i> JAP (Cincinnati)	Type B (Japanese)
6	FRAN025	<i>Francisella tularensis</i>	subsp. <i>tularensis</i> (VT68)	Type A1 (United States)
7	FRAN029	<i>Francisella tularensis</i>	subsp. <i>holarctica</i> (425)	Type B (United States)
8	FRAN031	<i>Francisella tularensis</i>	subsp. <i>tularensis</i> (Schemm)	Type A1 (United States)
9	FRAN072	<i>Francisella tularensis</i>	subsp. <i>tularensis</i> (WY96)	Type A2 (United States)
10	N/A	<i>Francisella tularensis</i>	Supsp. <i>mediasiatica</i>	

87
88
89
90
91

^a UCC = Department of Defense Unified Culture Collection; components available through Biodefense and Emerging Infections Research Resources Repository.

92
93
94

Table IV: Exclusivity Panel (near-neighbor)

No.	Species	Strain
1	<i>Francisella philomiragia</i>	Jensen O#319L ATCC 25015
2	<i>Francisella philomiragia</i>	Jensen O#319-029 ATCC 25016
3	<i>Francisella philomiragia</i>	Jensen O#319-036 ATCC 25017
4	<i>Francisella philomiragia</i>	Jensen O#319-067 ATCC 25018
5	<i>Francisella philomiragia</i>	D7533, GA012794
6	<i>Francisella philomiragia</i>	E9923, GA012801
7	<i>Francisella novicida</i>	D9876, GA993548
8	<i>Francisella novicida</i>	F6168, GA993549
9	<i>Francisella novicida</i>	U112, GA993550
10	<i>Francisella hispaniensis</i>	DSM 22475

120
121
122
123
124
125
126
127
128

129
130
131
132
133
134
135

Guidance

Organisms may be tested as isolated DNA, or combined to form a pool of isolated DNA. Isolated DNA may be combined into pools of up to 10 exclusivity panel organisms, with each panel organism represented at 10 times the AMDL, where possible. If an unexpected result occurs, each of the exclusivity organisms from a failed pool must be individually re-tested at 10 times the AMDL.

In silico screening shall be performed on signature sequences (e.g., oligo primers/probes/ amplicons) to predict specificity and inclusivity across all sequenced *Francisella* strains. *In silico* results are suggestive of potential performance issues. *Basic Local Alignment Search Tool* (BLAST) should be able to predict hybridization events between signature components and available *Francisella* genomic sequence data in GenBank®. Results of *in silico* analyses shall be included in method/assay performance evaluation reports.

136 **Annex 1: Environmental Factors For Validating Biological Threat Agent Detection Assays**

137

138 [Adapted from the Environmental Factors Panel approved by SPADA on June 10, 2010.]

139

140 The Environmental Factors Studies supplement the biological threat agent near-neighbor
141 exclusivity testing panel. There are three parts to Environmental Factors studies: part 1 -
142 environmental matrix samples; part 2 - the environmental organisms study; and part 3 - the
143 potential interferences applicable to Department of Defense applications.³

144

145

146 **Part 1:**

147

148 **Environmental Matrix Samples - Aerosol Environmental Matrices**

149

150 Method developers shall obtain environmental matrix samples that are representative and
151 consistent with the collection method that is anticipated to ultimately be used in the field. This
152 includes considerations that may be encountered when the collection system is deployed
153 operationally such as collection medium, duration of collection, diversity of geographical areas
154 that will be sampled, climatic/environmental conditions that may be encountered and seasonal
155 changes in the regions of deployment.

156

157 Justifications for the selected conditions that were used to generate the environmental matrix
158 and limitations of the validation based on those criteria must be documented.

159

- 160 • Method developers shall test the environmental matrix samples for interference using
161 samples inoculated with a target biological threat agent sufficient to achieve 95%
162 probability of detection.
- 163 • Cross-reactivity testing will include sufficient samples and replicates to ensure each
164 environmental condition is adequately represented.

165

166

³ Added in June 2015 for the Department of Defense project.

167

168 **Part 2: Environmental Panel Organisms** - This list is comprised of identified organisms from the
169 environment.

170

171 Inclusion of all environmental panel organisms is not a requirement if a method developer
172 provides appropriate justification that the intended use of the assay permits the exclusion of
173 specific panel organisms. Justification for exclusion of any environmental panel organism(s)
174 must be documented and submitted.

175

176 Organisms and cell lines may be tested as isolated DNA, or as pools of isolated DNA. Isolated
177 DNA may be combined into pools of up to 10 panel organisms, with each panel organism
178 represented at 10 times the AMDL, where possible. The combined DNA pools are tested in the
179 presence (at 2 times the AMDL) and absence of the target gene or gene fragment. If an
180 unexpected result occurs, each of the individual environmental organisms from a failed pool
181 must be individually re-tested at 10 times the AMDL with and without the target gene or gene
182 fragment at 2x the AMDL in the candidate method DNA elution buffer.

183

184 DNA in this list that already appear in the inclusivity or exclusivity panel do not need to be
185 tested again as part of the environmental factors panel.

186

187 • **Potential bacterial biothreat agents**

188 *Bacillus anthracis* Ames

189 *Yersinia pestis* Colorado-92

190 *Francisella tularensis* subsp. *tularensis* Schu-S4

191 *Burkholderia pseudomallei*

192 *Burkholderia mallei*

193 *Brucella melitensis*

194

195 • **Cultivable bacteria identified as being present in air soil or water**

196 *Acinetobacter lwoffii*

197 *Agrobacterium tumefaciens*

198 *Bacillus amyloliquefaciens*

199 *Bacillus cohnii*

200 *Bacillus psychrosaccharolyticus*

201 *Bacillus benzoovorans*

202 *Bacillus megaterium*

203 *Bacillus horikoshii*

204 *Bacillus macroides*

205 *Bacteroides fragilis*

206 *Burkholderia cepacia*

207 *Burkholderia gladioli*

208 *Burkholderia stabilis*

209 *Burkholderia plantarii*

210 *Chryseobacterium indologenes*

211 *Clostridium sardiniense*

212 *Clostridium perfringens*

213 *Deinococcus radiodurans*

214 *Delftia acidovorans*

215 *Escherichia coli* K12

216	<i>Fusobacterium nucleatum</i>
217	<i>Lactobacillus plantarum</i>
218	<i>Legionella pneumophila</i>
219	<i>Listeria monocytogenes</i>
220	<i>Moraxella nonliquefaciens</i>
221	<i>Mycobacterium smegmatis</i>
222	<i>Neisseria lactamica</i>
223	<i>Pseudomonas aeruginosa</i>
224	<i>Rhodobacter sphaeroides</i>
225	<i>Riemerella anatipestifer</i>
226	<i>Shewanella oneidensis</i>
227	<i>Staphylococcus aureus</i>
228	<i>Stenotrophomonas maltophilia</i>
229	<i>Streptococcus pneumoniae</i>
230	<i>Streptomyces coelicolor</i>
231	<i>Synechocystis</i>
232	<i>Vibrio cholerae</i>
233	
234	• Microbial eukaryotes
235	
236	<u>Freshwater amoebae</u>
237	<i>Acanthamoeba castellanii</i>
238	<i>Naegleria fowleri</i>
239	
240	<u>Fungi</u>
241	<i>Alternaria alternata</i>
242	<i>Aspergillus fumigatus</i>
243	<i>Aureobasidium pullulans</i>
244	<i>Cladosporium cladosporioides</i>
245	<i>Cladosporium sphaerospermum</i>
246	<i>Epicoccum nigrum</i>
247	<i>Eurotium amstelodami</i>
248	<i>Mucor racemosus</i>
249	<i>Paecilomyces variotii</i>
250	<i>Penicillium chrysogenum</i>
251	<i>Wallemia sebi</i>
252	
253	

254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298

- **DNA from higher eukaryotes**

- Plant Pollen⁴

- Zea mays* (corn)

- Pinus* spp . (pine)

- Gossypium* spp. (Cotton)

- Arthropods

- Aedes aegypti* (ATCC /CCL-125(tm) mosquito cell line)

- Aedes albopictus* (Mosquito C6/36 cell line)

- Dermatophagoides pteronyssinus* (Dust mite -commercial source)

- Xenopsylla cheopis* Flea (Rocky Mountain labs)

- Drosophila* cell line

- Musca domestica* (housefly) ARS, USDA, Fargo, ND

- Gypsy moth cell lines LED652Y cell line (baculovirus)– Invitrogen

- Cockroach (commercial source)

- Tick (*Amblyomma* and *Dermacentor* tick species for *F. tularensis* detection assays)⁵

- Vertebrates

- Mus musculus* (ATCC/HB-123) mouse

- Rattus norvegicus* (ATCC/CRL-1896) rat

- Canis familiaris*(ATCC/CCL-183) dog

- Felis catus* (ATCC/CRL-8727) cat

- Homo sapiens* (HeLa cell line ATCC/CCL-2) human

- Gallus gallus domesticus* (Chicken)

- Capri hirca* (Goat⁶)

- **Biological insecticides** – Strains of *B. thuringiensis* present in commercially available insecticides have been extensively used in hoaxes and are likely to be harvested in air collectors. For these reasons, it should be used to assess the specificity of these threat assays.

- B. thuringiensis* subsp. *israelensis*

- B. thuringiensis* subsp. *kurstaki*

- B. thuringiensis* subsp. *morrisoni*

- Serenade (Fungicide) *B. subtilis* (QST713)

Viral agents have also been used for insect control. Two representative products are:

Gypcheck for gypsy moths (*Lymanteria dispar* nuclear polyhedrosis virus)

Cyd-X for codling moths (Codling moth granulosis virus)

⁴ If pollen is unavailable, vegetative DNA is acceptable

⁵ Added by SPADA on (future approval date).

⁶ Added by SPADA on September 1, 2015.

299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319

Part 3: Potential Interferents Study

The Potential Interferents Study supplements the Environmental Factors Study, and is applicable to all biological threat agent detection assays for Department of Defense applications. Table 1a provides a list of potential interferents that are likely to be encountered in various Department of Defense applications.

Method developers and evaluators shall determine the most appropriate potential interferents for their application. Interferents shall be spiked at a final test concentration of 1 µg/ml directly into the sample collection buffer. 0 Sample collection buffers spiked with potential interferents shall be inoculated at 2 times the AMDL (or AMIL) with one of the target biological threat agents.

Spiked / inoculated sample collection buffers shall be tested using the procedure specified by the candidate method. A candidate method that fails at the 1 microgram per ml level may be reevaluated at lower concentrations until the inhibition level is determined.

It is expected that all samples are correctly identified as positive.

320
321

Table 1a: Potential Interferents

Compounds		Potential Theaters of Operation
group 1: petroleum-based	JP-8 ¹	airfield
	JP-5 ²	naval
	diesel/gasoline mixture	ground
	fog oil (standard grade fuel number 2)	naval, ground
	burning rubber ³	ground, airfield
group 2: exhaust	gasoline exhaust	ground
	jet exhaust	naval, airfield
	diesel exhaust	ground
group 3: obscurants	terephthalic acid ⁴	ground
	zinc chloride smoke ⁵	ground
	solvent yellow 33 ⁶	ground
group 4: environmental	burning vegetation	ground, airfield
	road dust	ground
	sea water (sea spray)	naval
group 5: chemicals	brake fluid ⁷	all
	brake dust ⁸	ground
	cleaning solvent, MIL-L-63460 ⁹	all
	explosive residues a) high explosives ¹⁰ b) artillery propellant ¹¹	all

322
323
324

325
326
327
328


Table 1a is offered for guidance and there are no mandatory minimum requirements for the number of potential interferents to be tested.

¹ **JP-8.** Air Force formulation jet fuel.

² **JP-5.** A yellow kerosene-based jet fuel with a lower flash point developed for use in aircraft stationed aboard aircraft carriers, where the risk from fire is particularly great. JP-5 is a complex mixture of hydrocarbons, containing alkanes, naphthenes, and aromatic hydrocarbons.


³ **Burning rubber** (tire smoke). Gaseous C1-C5 hydrocarbons: methane; ethane; isopropene; butadiene; propane. Polycyclic aromatic hydrocarbons (58-6800 ng/m³): parabenzo(a)pyrene; polychlorinated dibenzo-p-dioxins (PCDD); polychlorinated dibenzofurans (PCDF). Metals (0.7 - 8 mg/m³): zinc; lead; cadmium.

⁴ **Terephthalic acid.** Used in the AN/M83 hand grenade currently used by US military.



⁵ **Zinc chloride smoke.** Also known as “zinc chloride smoke” and “HC smoke”. Was used in the M8 grenade and still used in 155mm artillery shells. HC smoke is composed of 45% hexachloroethane, 45% zinc oxide, and 10% aluminum.

⁶ **Solvent yellow 33** [IUPAC name: 2-(2-quinoly)-1,3-indandione] is a new formulation being developed for the M18 grenade.



⁷ **Brake fluid.** DOT 4 is the most common brake fluid, primarily composed of glycol and borate esters. DOT 5 is silicone-based brake fluid. The main difference is that DOT 4 is hygroscopic whereas DOT 5 is hydrophobic. DOT 5 is often used in military vehicles because it is more stable over time requires less maintenance

⁸ **Brake dust.** Fe particles caused by abrasion of the cast iron brake rotor by the pad and secondly fibers from the semi metallic elements of the brake pad. The remainder of the dust residue is carbon content within the brake pad.

⁹ **MIL-L-63460,** "Military Specification, Lubricant, Cleaner and Preservative for Weapons and Weapons Systems"; trade name “Break-Free CLP”. Hyperlink: [Midway USA](#).

¹⁰ **High explosives.** The M795 155mm projectile is the US Army / Marine Corp's current standard projectile containing 10.8 kg of TNT. The M795 projectile replaced the M107 projectile that contained Composition B which is a 60/40 mixture of RDX/TNT. RDX is cyclotrimethylene trinitramine. Suggestion: test RDX/TNT together.

¹¹ **Artillery propellant.** Modern gun propellants are divided into three classes: single-base propellants which are mainly or entirely nitrocellulose based, double-base propellants composed of a combination of nitrocellulose and nitroglycerin, and triple base composed of a combination of nitrocellulose and nitroglycerin and nitroguanidine. Suggestion: test total nitrocellulose/ nitroglycerin nitroguanidine together.



AOAC Stakeholder Panel on Agent Detection Assays (SPADA)

AOAC STANDARDS DEVELOPMENT PROCESS

Stakeholder Panel Working Groups


Deborah McKenzie
Sr. Director,
AOAC Standards Development
AOAC INTERNATIONAL
September 1, 2015

AOAC INTERNATIONAL HEADQUARTERS
2275 Research Blvd, Ste 300
Bethesda, Maryland 20814



The Science® Association Dedicated to Analytical Excellence®

AOAC Standards Development Processes




AOAC INTERNATIONAL

As an international standards development organization, AOAC must maintain the following principles throughout all its standard setting activities:

- Transparency
- Openness
- Balance of Interests
- Due Process
- Consensus
- Appeals



Stakeholder Panel Role and Output



AOAC Voluntary Consensus Standards

- **Standard Method Performance Requirements (SMPRs)**
 - Published in *Official Methods of Analysis of AOAC INTERNATIONAL*
 - Manuscript published in *Journal of AOAC INTERNATIONAL*



Stakeholder Panel Working Groups

- Engage in the detailed discussions and work of the stakeholders
- Develop draft fitness for purpose and standard method performance requirements (SMPRs) or other draft standard as proposed by stakeholder panel
- Recommend draft standards to the stakeholder panel
- Can review and recommend methods for consideration
- Formed based on efficiency for work
- Managed by staff
- Conducted in accordance with national and international standards development criteria



Working Group Membership

- Working group membership is not vetted
- Membership is a microcosm of stakeholder panel
 - Experts from government, industry, research, states, military, public health sector, public safety sector, method developers, method end users, etc...
- Moderated by working group chair(s)
- Facilitated by AOAC staff
- Working groups are often formed during the stakeholder meetings and joint working group meetings



Working Group Chair(s)

- Make a launch presentation to the stakeholder panel introducing the topic of the standard
- Moderate the working group discussions
- Work with AOAC CSO and staff
 - On presentations
 - Draft standard
 - Reconcile comments on standard
- Make a SMPR® presentation proposing the draft SMPR to the stakeholder panel.



Working Group Meetings

- In person meetings
 - Wednesday, September 2, 2015
- Teleconference meetings
 - To be scheduled after SPADA meetings
- Facilitated by AOAC staff
- Moderated by Working Group chair(s)
- Discuss the contents of SMPR®
- Goal of Working Group deliberations
 - To reach a general agreement on draft SMPR®



Draft Standards

- AOAC will post the draft SMPR® for a 30 day public comment period
- AOAC CSO and Working Group Chair will reconcile comments
 - Revise SMPR® if needed
 - Reconvene working group if needed



Standard Methods Performance Requirements (SMPRs)

Standard Methods Performance Requirements

AOAC SMPR 011.006

Standard Methods Performance Requirements for Fatigue in Infant Formula and Adult Pediatric Nutritional Formula

Approved by: *Standardization Panel on Infant Formula and Adult Pediatric Nutrition*

Final Review Date: April 8, 2011

Effective Date: April 8, 2011

Method Use:

1. **Application:**

2. **Reference:**

3. **Method Performance:**

4. **Method:**

5. **Method Acceptance:**

6. **Method Revision:**

7. **Method Approval:**

8. **Method Approval:**

9. **Method Approval:**

10. **Method Approval:**

11. **Method Approval:**

12. **Method Approval:**

13. **Method Approval:**

14. **Method Approval:**

15. **Method Approval:**

16. **Method Approval:**

17. **Method Approval:**

18. **Method Approval:**

19. **Method Approval:**

20. **Method Approval:**

21. **Method Approval:**

22. **Method Approval:**

23. **Method Approval:**

24. **Method Approval:**

25. **Method Approval:**

26. **Method Approval:**

27. **Method Approval:**

28. **Method Approval:**

29. **Method Approval:**

30. **Method Approval:**

31. **Method Approval:**

32. **Method Approval:**

33. **Method Approval:**

34. **Method Approval:**

35. **Method Approval:**

36. **Method Approval:**

37. **Method Approval:**

38. **Method Approval:**

39. **Method Approval:**

40. **Method Approval:**

41. **Method Approval:**

42. **Method Approval:**

43. **Method Approval:**

44. **Method Approval:**

45. **Method Approval:**

46. **Method Approval:**

47. **Method Approval:**

48. **Method Approval:**

49. **Method Approval:**

50. **Method Approval:**

51. **Method Approval:**

52. **Method Approval:**

53. **Method Approval:**

54. **Method Approval:**

55. **Method Approval:**

56. **Method Approval:**

57. **Method Approval:**

58. **Method Approval:**

59. **Method Approval:**

60. **Method Approval:**

61. **Method Approval:**

62. **Method Approval:**

63. **Method Approval:**

64. **Method Approval:**

65. **Method Approval:**

66. **Method Approval:**

67. **Method Approval:**

68. **Method Approval:**

69. **Method Approval:**

70. **Method Approval:**

71. **Method Approval:**

72. **Method Approval:**

73. **Method Approval:**

74. **Method Approval:**

75. **Method Approval:**

76. **Method Approval:**

77. **Method Approval:**

78. **Method Approval:**

79. **Method Approval:**

80. **Method Approval:**

81. **Method Approval:**

82. **Method Approval:**

83. **Method Approval:**

84. **Method Approval:**

85. **Method Approval:**

86. **Method Approval:**

87. **Method Approval:**

88. **Method Approval:**

89. **Method Approval:**

90. **Method Approval:**

91. **Method Approval:**

92. **Method Approval:**

93. **Method Approval:**

94. **Method Approval:**

95. **Method Approval:**

96. **Method Approval:**

97. **Method Approval:**

98. **Method Approval:**

99. **Method Approval:**

100. **Method Approval:**

- Documents a community's analytical method needs.
- Very detailed description of the analytical requirements.
- Includes method acceptance requirements.
- Used to adopt AOAC Official Methods by Expert Review Panels.
- Published as a standard.



Documentation and Communication

- AOAC carefully documents the actions of Stakeholder Panel and the Working Groups
- AOAC will prepare summaries of the meetings
 - Communicate summaries to the stakeholders
 - Publish summaries in the *Referee* section of AOAC's *Inside Laboratory Management*
- AOAC publishes its voluntary consensus standards and *Official Methods*
 - *Official Methods of Analysis of AOAC INTERNATIONAL*
 - *Journal of AOAC INTERNATIONAL*
- AOAC publishes the status of standards in the Referee section of AOAC's *Inside Laboratory Management*



Working Groups - Timelines

- **September 1, 2015:**
 - Launching of new working group topic of new standards
- **September 2, 2015:**
 - First meeting of the working groups
- **September through December 2015:**
 - Teleconferences of working groups
- **January – February 2016:**
 - Public Comment Period
- **February – March 2016:**
 - Reconciliation of Comments
- **March 2016:**
 - Presentation of draft SMPR during next SPADA meeting



SPADA Chair & Working Group Chairs

***B. anthracis* Working Group (BAWG)**

Paul Jackson (LLNL/LANL) & Ted Hadfield (Hadeco)

***Y. pestis* Working Group (YPWG)**

Luther Lindler (DHS)

***F. tularensis* Working Group (FTWG)**

Peter Emanuel (DoD) & Mark Wolcott (DoD);
Paul Keim (NAU) & David Wagner (NAU)

Environmental Factors Working Group (EFWG)

Stephen Morse (CDC)

Public Health Actionable Assay Working Group (PHAAWG)

Peter Estacio (LLNL)

***B. anthracis* Handheld Assay Working Group (BaHHAWG)**

Marian McKee (BioReliance Corp.)

Ricin Handheld Assay Working Group (RicinHHAWG)

Mark Poli (DoD)

***Burkholderia* Working Group (BurkWG)**

Paul Keim (NAU) & Alex Hoffmaster (CDC)

Assay Controls Working Group (ACWG)

Christina Egan (NYS DH) & Larry Blynn (Ibis)

Variola Working Group (VWG)

Victoria Olson (CDC) & Ted Hadfield (Hadeco)

***Coxiella burnetii* (CWG)**

Linda Beck (NSWC) & James Samuels (Texas A&M)

***Staphylococcus* Enterotoxins (SEWG)**

Sandra Tallent (US FDA)

Venezuelan Equine Encephalitis (VEEWG)

Eileen Ostlund (USDA)



13

Contact Information

Contact AOAC Staff:


Tel: 301.924.7077

Web: www.aoac.org





- **E. James Bradford**, Executive Director & CEO, jbradford@aoac.org, ext. 102
- **Krystyna McIver**, Executive for Scientific Engagement and Communication, kmciver@aoac.org, ext. 111
- **Scott Coates**, Chief Scientific Officer, scoates@aoac.org, ext. 137
- **Deborah McKenzie**, Sr. Director – Standards Development and AOAC Research Institute, dmckenzie@aoac.org, ext. 157




Assay for *C. Botulinum* toxins



Seven Serotype, A - G






Toxin Complex



U.S. Food and Drug Administration
Protecting and Promoting Public Health

www.fda.gov

Toxin Complex 500 kDA



Detection

Structural element

- Relatively easy to detect
- Presence or absence
- Often qualitative
- inexpensive

Functional

- Relatively Not-easy to detect
- Active or inactive
- Qualitative and quantitative
- Expensive
- Representative of BoNT intoxication

Types of testing


Mouse assay
(detection of functional toxin)

toxin screening


toxin titer

toxin neutralization
by using antitoxin

Animal requirements



Alternate methods





Detection techniques for botulinum neurotoxins

Method	Limit of Detection (ml ⁻¹)	Multiplex Format	Sample-Matrix Interference	Automation
1. Mouse Bioassay *	20 pg	No	Limited	No
2. ELISA*	5 pg - 2 ng	Limited	Manageable	Limited
3. ECL*	5 pg -50 ng	Limited	Manageable	Limited
4. Lateral flow assay*	5-50 ng	No	High	N/A
5. Column flow assay*	1-50 ng	No	High	N/A
6. Flow cytometry assay	50 pg-20 ng	Yes	Manageable	Yes
7. Immuno-PCR	1 pg- 5 pg	Limited	Manageable	No
8. L-PCR	0.02 fg	Limited	N/A	No
9. BDG assay	100 ng	Yes	Low	Yes
10. Array biosensor assay	40-200 ng	Yes	Low	Yes
11. Aptamer electrochemical assay	40 pg	Yes	Low	Yes
12. Peptide array assay	3 pg	Yes	High	Yes
13. ALISSA	0.5 fg	No	Low	No
14. Endopep-MS assay*	0.4 -6 pg	Yes	High	Yes
15. Cell based assays	1-10 ng	No	High	No

(* assay used for food analysis); Enzyme-linked immunosorbent assay (ELISA); Electro-chemiluminescent (ECL) assay; Liposomal-PCR (L-PCR); Bidiffractive grating (BDG) assay; Assay with a large immunosorbent surface area (ALISSA)



Fitness for Purpose

To validate a rapid multi-plex *Clostridium botulinum* neurotoxin detection assay that has the ability to:

- Detect and differentiate serotypes A through G
- Detect bivalent toxins effectively and also determine the dominant type
- Differentiate complex toxin with associated proteins from purified toxin (di-chain) and toxoid



The Scientific Association Dedicated
to Analytical Excellence[®]

STAKEHOLDER PANEL ON AGENT DETECTION ASSAYS (SPADA)

Background & Fitness for Purpose

Brucella suis

Frank F. Roberto, PhD
Rockville, MD
23 March 2016



The Scientific Association Dedicated
to Analytical Excellence[®]

Background

- Brucellosis, aka Malta Fever, undulant fever, Bang's Disease, fistulous withers (in horses)
- The causal agent is an intracellular bacterial pathogen, first isolated after cases of the disease in garrison troops on the island of Malta who drank infected goat's milk (Sir David Bruce, "Micrococcus melitensis")
- Pasteurization of dairy products and culling of infected animals have reduced the prevalence of the disease in many domesticated species, but wildlife reservoirs remain throughout the world including US



The Scientific Association Dedicated
to Analytical Excellence[®]

Background

- *B. suis* anecdotally first pathogen developed as a biological weapon by former US offensive program (W. Patrick, III, personal communication)
- Infectious dose of 10-100 organisms validated in historic “8-ball” chamber at Ft. Detrick with human subjects
- Aerosol transmission (first noted in swine abattoirs), but also via fomites and sexual route documented in animals and humans
- Serious acute and chronic symptoms of brucellosis often complicate correct diagnosis



The Scientific Association Dedicated
to Analytical Excellence[®]

Background

- α -proteobacteria, Order Rhizobiales, Family Brucellaceae
- Other Brucellaceae include Mycoplana, Ochrobactrum, Pseudochrobactrum, Paenochrobactrum, and Crabtreeella.
- Ten currently recognized species (often with strong host-specificity) further divided into biovars
- Members of the Rhizobiales are similar enough to generate false positives (eg *Agrobacterium tumefaciens*)




Background

- Confirmation by isolation remains the gold standard, in spite of aerosol hazard and low infectious dose
- Cultivation often includes antibiotics and antibiotic dyes (thionin, basic fuchsin) to counterselect against other microbes
- Select agent regulations complicate this process! Proposed rule changes out for comment



Genome-based *Brucella* taxonomy


Bohlin et al., 2010. BMC Evol. Biol. 10:249





The Scientific Association Dedicated
to Analytical Excellence[®]

Remarkable amount of gene synteny




Wattam et al., 2009 *J Bacteriol* 191



The Scientific Association Dedicated
to Analytical Excellence[®]

National Prevalence Rate: Brucellosis Affected Cattle Herds

FY 2004 = 0.0005% FY 2006 = 0.0002% FY 2008 = 0.0003%
FY 2005 = 0.0003% FY 2007 = 0.0001% FY 2009 = 0.0000%
FY 2010 = 0.0001% (as of August 1, 2010)






The Scientific Association Dedicated
to Analytical Excellence[®]

Brucellosis in the Greater Yellowstone Area

- What is the real potential for interspecies transmission between large ungulates and cattle in the Intermountain West?
- Can rapid, accurate molecular diagnostics improve management and minimize impacts to wildlife?



The Scientific Association Dedicated
to Analytical Excellence[®]

Real-time PCR assay for *B. abortus*

Newby et al. 2003. *Appl. Environ. Microbiol.* 69



- *B. abortus* specific (tested against panel of over 100 strains)
- 7.5 fg limit of detection (ca. 2 genomic copies)
- Semi-quantitative nature of real-time PCR permits estimation of bacterial load in samples
- Detection in 15-30 minutes
- No sample prep necessary in some cases
- Hybridization probes allow discrimination of amplicons based on post-amplification melt curves (potential to identify S19 and RB51 without multiplexing)
- New TaqMan assay developed for other instruments (15 fg LOD)



The Scientific Association Dedicated to Analytical Excellence[®]

Brucellosis in the Greater Yellowstone Area

- 600-900 animals will be lethally removed this spring – again (record 1726 in 2008)
- Can science turn the tide?



The Scientific Association Dedicated to Analytical Excellence[®]

Clear evidence that elk pose a greater risk to cattle in the GYA

Rhyhan, et al. 2013 *EID* 19

- Cattle
- Bison, ranched
- Elk, wild
- Bison, wild




Table. Cattle and ranched bison herds found infected with *Brucella abortus* due to transmission from elk, Greater Yellowstone Area, USA

Herd no.	County, state	Species	Herd size	Date detected	Seropositive, %	Culture results	Distance to feeding ground, km
1	Fremont, ID	Cattle	50-100	2002 Apr	12.0	Biovar 1	50*
2	Sublette, WY	Cattle	>300	2003 Oct	9.9	Biovar 1	2.4
3†	Teton, WY	Cattle	>300	2004 Jun	1.9	Biovar 4	Adjacent
4	Bonneville, ID	Cattle	<50	2005 Aug	20.0	Biovar 1	85‡
5	Park‡, MT	Cattle	>300	2007 May	0.2	Biovar 1	>100
6	Park, MT	Cattle	<50	2008 May	2.9	Biovar 1	>100
7	Sublette, WY	Cattle	>300	2008 Jun	5.5	Biovar 4	24
8	Jefferson, ID	Cattle	>300	2009 Jul	1.5	Biovar 1	85
9	Park, WY	Cattle	>300	2010 Oct	1.1	Biovar 1	>100
10	Park, WY	Bison	200-300	2010 Nov	11.5	Biovar 4	>100
12	Park, WY	Cattle	>300	2011 Feb	0.9	Biovar 1	>100
13	Park, WY	Cattle	>300	2011 Sep	1.2	Biovar 1	>100
14	Park, MT	Cattle	>300	2011 Sep	2.0	Biovar 1	>100
15	Madison, MT	Bison	>300	2011 Nov	0.2	Biovar 1	>100
16	Fremont, ID	Cattle	50-100	2012 Apr	5.8	Biovar 1	90
17	Bonneville, ID	Bison	200-300	2012 Mar	0.7	Biovar 4	40

*Elk were also intentionally fed on ranch by owner.

†Brucellosis was detected in 2 herds that were pastured together during spring of 2004. This was considered a single transmission event, and statistics are given for the combined herds.


‡Herd located 0.8 km from site where elk feeding ground had been until 2003.

§This herd was discovered infected in Carbon County, MT, in 2007, but animals had been transported from Park County in 2005. Index cow aborted in 2005 and did not calve in 2006.



The Scientific Association Dedicated
to Analytical Excellence[®]

Increase in seropositive feral swine




Leiser et al., 2013. *Vet Microbiol* 166



The Scientific Association Dedicated
to Analytical Excellence[®]

Genomic alignments can inform assays

Real-time PCR assay development




Roberto et al. 2009 *Unpublished* (DHS)



The Scientific Association Dedicated
to Analytical Excellence[®]

Real-time PCR assay for *B. suis*



- *B. suis* bv.1 (may also detect 2-4, but results inconclusive with only one strain of each available)
- 2 fg limit of detection (less than 1 genomic copy)
- TaqMan assay developed on ABI 7000 and 7900HT Fast block (25µl rxn)
- Detection in 15-30 minutes

Roberto et al. 2009 *Unpub* (DHS)



The Scientific Association Dedicated
to Analytical Excellence[®]

Proposed Changes to FSAP Status

- Removal of *B. abortus*, *B. melitensis*, and *B. suis* from HHS list of overlap agents because (1) mortality rate is low; (2) brucellosis is treatable with antibiotics; and (3) human-to-human transmission is “extremely rare” and zoonotic transfer is also rare (each point cites Lecaroz et al., 2006, *J. Antimicrob. Chemotherap.* 58:549-546) – in spite of at least 3 human cases associated with feral swine hunting; dogs at risk
- USDA/APHIS proposes to remove *B. abortus* and *B. suis*



The Scientific Association Dedicated
to Analytical Excellence[®]

Significance

- Is brucellosis truly eradicated in the United States?
- Is *B. suis* distinguishable from *B. canis*?
- Do feral swine pose a greater risk to human and animal health than Yellowstone bison and elk?
- Does our working group feel we can address the standard method performance requirements for *B. suis* real-time PCR assays, as well as for *B. abortus* and *B. melitensis*?



The Scientific Association Dedicated
to Analytical Excellence[®]

General Analytical Needs

- Working group tasked with addressing *B. suis*
- Working group to decide if *B. abortus* and *B. melitensis* can be addressed at the same time
- Detection in aerosols (air collection of particulates onto filters)
- Inclusion panels - tbd
- Exclusion panels – tbd, but also a common SMPR panel that has been established from air sampling experiences (eg BioWatch)
- Chemical contaminant background



The Scientific Association Dedicated
to Analytical Excellence[®]

Challenges - 10 species currently described



Many demonstrate host specificity



The Scientific Association Dedicated
to Analytical Excellence[®]

Existing Methods

- Foster et al. 2008 SNP-based real-time PCR assays for all major Brucella clades
- Fretin et al. 2008 SNP-based real-time PCR assays for B. suis (137 field strains)
- Hinić et al. 2008 Conventional and real-time PCR assays for multiple Brucella species
- Lopez-Goñi et al. 2016 Multiplex “Bruce-ladder” for all known species
- Scholl et al., 2010 Multiplex PCR assay for all described species
- Yu and Nielsen, 2010 Croatian Med J 51 review



The Scientific Association Dedicated
to Analytical Excellence[®]

Fitness for Purpose (proposal)

Real-time PCR assay that unambiguously identifies *Brucella suis* (all biovars?) from *B. canis* and other currently-described species and biovars of *Brucella* with single genomic copy sensitivity (fg) in 4 hours or less. The assay must not generate false positives in a background that contains closely-related genera such as *Agrobacterium*, *Rhizobium* and *Ochrobactrum*.



The Scientific Association Dedicated
to Analytical Excellence[®]

QUESTIONS or COMMENTS?



The Scientific Association Dedicated
to Analytical Excellence[®]

STAKEHOLDER PANEL ON AGENT DETECTION ASSAYS (SPADA)

Background & Fitness for Purpose

Burkholderia pseudomallei

Jay E. Gee, PhD

Centers for Disease Control and Prevention

Rockville, MD


March 23, 2016



The Scientific Association Dedicated
to Analytical Excellence[®]

Burkholderia pseudomallei

- Tier 1 Select Agent
- Causes disease melioidosis
- Aerobic, gram-negative bacillus
- Naturally in water and soil in endemic areas
- Opportunistic pathogen
- Causes human and animal disease
- Incubation period: 1 to 21 days, mean of 9 days
- Can be latent for decades before causing illness





The Scientific Association Dedicated
to Analytical Excellence[®]

Routes of Infection

- Inhalation: aerosolized in water or soil
- Inoculation: skin wound contact with soil or water
- Ingestion: water
- Person-to-person: rare
- Animal-to-human: not observed



<http://www.vhfcn.org/unitlinks.html>



<http://www.ndm.ox.ac.uk/principal-investigators/researcher/david-dance>



The Scientific Association Dedicated
to Analytical Excellence[®]

Symptoms

- “The Great Mimicker”
- Acute pulmonary infection
 - most common presentation
- Localized infection
- Disseminated infection
- Fulminant septicemia
- Neurological (less common)




CHEST 2011; 140(1):239-242



The Scientific Association Dedicated
to Analytical Excellence[®]

Geography




<http://www.cdc.gov/melioidosis/prevention/>



The Scientific Association Dedicated
to Analytical Excellence[®]

Geography


From: Predicted global distribution of *Burkholderia pseudomallei* and burden of melioidosis



Areas of high environmental suitability are shown in red and areas of low suitability in green.



Genetics



<http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2784454/>



Significance

- Improvements in rapid detection are desirable
- A uniform panel of isolates will allow standardization to better compare analytical test results



The Scientific Association Dedicated
to Analytical Excellence[®]

General Analytical Needs

- Inclusivity Panel for *B. pseudomallei*
 - Genetic diversity
 - Geographic diversity
 - Available from accessible archives
- Exclusivity Panel
 - Closely related to *B. pseudomallei*
 - May cause diagnostic confusion



The Scientific Association Dedicated
to Analytical Excellence[®]

General Analytical Needs

- Define acceptable:
 - limit of detection
 - Sensitivity
 - Specificity



*The Scientific Association Dedicated
to Analytical Excellence[®]*

Challenges

- Infectious dose for humans – unknown
- Sampling bias – strains in archives predominantly from Southeast Asia and Australia



*The Scientific Association Dedicated
to Analytical Excellence[®]*

Fitness for Purpose (proposal)

- Motion to compose list of bacteria suitable for inclusivity and exclusivity panels
- Motion to develop acceptable analytical standards for:
 - Limit of Detection
 - Sensitivity
 - Specificity



*The Scientific Association Dedicated
to Analytical Excellence[®]*

QUESTIONS?



Smallpox: Considerations for testing


Victoria A. Olson Ph.D.
Poxvirus and Rabies Branch
Centers for Disease Control and Prevention

March 22, 2016

National Center for Emerging and Zoonotic Infectious Diseases
Division of High Consequence Pathogens and Pathology



Poxvirus 101



Family of large, double stranded DNA viruses

- Within genera, antigenic similarity: cross protection

Complex viruses, cytoplasmic lifecycle

Genus *Orthopoxvirus*:

- 90-98% nucleotide identity across species
- *Variola*, *Vaccinia*, *Cowpox*, *Monkeypox* – all can cause human disease
 - *Variola* evolved to be a sole human pathogen: SMALLPOX
- *Camelpox*, *Ectromelia*, *Taterapox* are NOT known to cause human disease

Objectives

- Smallpox introduction
- Development of smallpox clinical diagnostics
- Developing a framework for considering environmental smallpox detection/testing
 - Results of 2014 Standard Method Performance Requirements for Detection and Identification of *Variola virus* DNA in Aerosol Collection Filters and/or Liquids

Smallpox disease elimination and eradication
Variola virus elimination and (?) eradication


SMALLPOX HISTORY AND BACKGROUND



Smallpox

Disease

- ❑ **Sole human pathogen**
- ❑ **No known non-human animal reservoir**
- ❑ **Transmitted by respiratory route (largely airborne droplets)**
 - Rare, but notable occurrences of airborne transmission in some hospitals
- ❑ **Transmitted by percutaneous exposure**
- ❑ **Fomites – rare cause of transmission**
- ❑ **Not foodborne or waterborne**



Progression of smallpox

- Incubation Period
- Pre-eruptive Stage:
 - abrupt onset high fever/constitutional symptoms
- Macules
- Papules
- Vesicles
- Pustules
- Scabs
- Scars



Smallpox - disease elimination/eradication

- ❑ **Disease – viral exanthem**
 - Major (“avg” 30% CFR) and minor disease (<1% CFR)
- ❑ **Disease prevention**
 - Childhood vaccination – variable rates; +/-variolation
- ❑ **Global Smallpox Eradication program - 1958**
 - Intensification of Smallpox eradication program 1967
 - Surveillance -> contact tracing, vaccination of contacts (and contacts)
 - Isolation of cases, observation of contacts
 - Lyophilized vaccine, semi-standardized vaccine production
 - *Vaccinia virus*
 - “Take” ~ protection
 - No non-human animal reservoir
- ❑ **Elimination in all countries by 1977**
 - Commission to Certify Smallpox eradication activities
 - Certified as eradicated in 1979
 - WHA : Declaration of smallpox eradication 1980

* Smallpox and its Eradication WHO 1988

History- *Variola virus* elimination and eradication

Consolidation of laboratory-held virus materials*

- 1975 survey by WHO, post lab exposure in 1973 (LSTMH)
 - 74 labs report *Variola virus* materials
- 1976 voluntary consolidation
 - 1978 – Birmingham, England smallpox “lab”: 1 death, illness
- 1979 – WHO Committee of Experts recommends to preserve *Variola virus* stocks in a few collaborating center (CC) labs, review in 1982:->19 recommendations by the Global Commission
 - 1979 – 7 labs report *Variola virus* stocks
 - 1981 – 4 CC labs with *Variola virus*
 - Periodic inspections for safe and secure use virus
- 1984 – consolidation of stocks to 2 WHO Collaborating Centers – BSL-4 facilities (WHA 33.4)

* Smallpox and its Eradication 1988

Virus eradication – considerations to 1999

- ❑ **Additional Global Commission sanctioned research – reflect (new) technologies of the time**
 - Cloning of *Variola virus* genomes – in representative segments
 - Hybrid viruses* (1981): proof of recombination/“transfection”
 - Scientific Advisory Group of Experts (1984)
 - Vaccine research using *Vaccinia virus* vector
 - Sequencing of virus genomes –
 - 1993 - Two complete *Variola virus* “major” genomes available
- ❑ **Bioterrorism threat once vaccination program ceases**
- ❑ **Decision to prohibit genetic manipulation of *Variola virus*, restrict access to genomic elements and genome**
- ❑ **Reports that Russia had attempted to “weaponize” *Variola virus***

* Sam and Dumbell Expression of poxvirus DNA in coinfecting cells and marker rescue of thermosensitive mutants by subgenomic fragments of DNA *Annales de virologie*, 1981

WHO/WHA process
IOM 1999, 2009
research 2000 to today

SMALLPOX RESEARCH AGENDA: DIAGNOSTICS

Smallpox Research Agenda: focused on preparedness needs

- Institute of Medicine Report recommendations for “Assessment of Future Scientific Needs for Live *Variola Virus*” (1999) have helped to frame the research agenda.
 - Protocols approved by WHO technical subcommittee
 - Research updates provided annually to the WHO Advisory Committee for *Variola Virus* Research
 - Collaborative HHS (largely CDC) and DoD (largely USAMRIID)
- All U.S. work with live *Variola virus* occurs within the BSL-4 containment laboratory at the CDC
 - Inspected regularly by U.S. security and biosafety authorities and WHO biosafety teams
 - Genetic manipulation of *Variola virus* not authorized by WHO
 - 1994 Ad hoc *Orthopoxvirus* Advisory Committee recommendation
 - **Full genomes of *Variola virus* can only be maintained at the 2 WHO CCs**
 - **No lab can have more than 20% of the *Variola virus* genome, except a WHO Collaborating Center**
- All research findings to be made available to the international scientific community

IOM recommendations* 1999-> WHO sanctioned research agenda

- Molecular characterization of *Variola virus* for more sensitive and specific diagnostic development
 - Sequencing entire genomes and specific genes
- Antiviral
- Less reactogenic vaccine development
- Animal model – pathogenesis, model system for antiviral & novel vaccine evaluation
- *Fundamental research – host pathogen interaction*

* Assessment of Future Scientific Needs for Live *Variola Virus*; N.A.Press (1999)

WHA resolutions and WHO Protocol approval process

- 1999 WHA resolution - postpone decision on destruction until 2002
- 2002 WHA resolution - postpone decision on destruction
- 2005 WHA
 - Increased focus on “essential” public health research
 - Interpreted by WHO committee to preclude fundamental research
 - Major review of the research to the WHA in 2011
 - Advisory Group of Independent Experts (AGIES) review *Variola virus* research in 2010
- 2011 WHA
 - Resolution to revisit in 3 years
 - AGIES conduct second review of *Variola virus* research in 2013
- 2013 WHA
 - Request to consider question of synthetic biology
 - Report to be shared 2016


Virus characterization

“Genomic sequencing and limited study of *Variola virus* surface proteins derived from geographically dispersed specimens is an essential foundation for important future work. Such research could be carried out now, and could require a delay in the destruction of known stocks, but would not necessitate their indefinite retention.*”

* Assessment of Future Scientific Needs for Live *Variola Virus*; N.A.Press (1999)

Maximum likelihood analysis of single nucleotide matrices (*Variola virus*)

48 isolates passaged
2-3 times to
high titer master
seed
stocks
DNA extracted
Sequenced using
primer walking
methodology



Li Y. et al. PNAS 2007;104:15787-15792 Esposito et al. Science 2006

Diversity of *Variola virus* ~200 kb dsDNA, ~200 ORFs

- Diversity of *Variola virus* strains is associated with geographic distance
- Alastrim minor (South America) / Variola major (Asia): ~600 SNPs, ~80 Indels
- Alastrim minor / Variola intermediate (West Africa): ~350 SNPs, ~45 Indels
- Variola African minor/major / Variola major (Asia): ~150 SNPs, ~30 Indels



- Central region: virion structural proteins, enzymes - 30 gene sequences are perfectly conserved or have only synonymous SNPs, highly conserved, essential function.
- Left and Right end regions: Host range and immunomodulatory genes - majority of Indels/frameshift mutations, fragmented sequences, additional/absent of ORFs, - likely reflecting selection pressures.
- **Versus other Orthopoxviruses:**
- *Variola / Camelpox-Taterapox viruses:* ~3200 SNPs, ~380 Indels
- *Variola / Monkeypox virus:* ~7500 SNPs, ~600 Indels

Diagnosics/environmental detection

“If further development of procedures for the environmental detection of *Variola virus* or for diagnostic purposes were to be pursued, more extensive knowledge of the genome variability, predicted protein sequences, virion surface structure, and functionality of *Variola virus* from widely dispersed geographic sources would be needed.*”

* Assessment of Future Scientific Needs for Live *Variola Virus*; N.A.Press (1999)

Diagnosics

- Why “if?”
- Proponents, in 1999, that EM and standard PCR techniques were sufficient for smallpox diagnostics
 - As of 2002 – survey of EM capacity in state health departments reveals only 3-8 with skilled capability to any capability
 - Newer technology: real time PCR

Diagnosics: Nucleic acid testing Real time PCR Assays

- Platform supported at Laboratory Response Network (LRN)
- High throughput
- Sensitive/specific
 - Sensitive to 1-50 genome copies
 - Historically lesion samples contained 10^4 - 10^7 infectious virions
 - Assays validated against authentic *Variola virus* genomic material
- Limitations:
 - time to get samples to reference labs

Diagnosics developed


- CDC developed/evaluated (real time) PCR assays targeting *Orthopoxvirus* genus and various species (*Variola*, *Monkeypox*, *Vaccinia*, etc.)
 - Provide reagents/facilities for others to evaluate assays
 - ~Thirteen peer-reviewed publications evaluate PCR assays against authentic *Variola virus* genomic material
- Subset used in LRN (*Variola*, *Monkeypox*, *Cowpox* and *Vaccinia virus* detection)
 - 2002 onward: vaccine AE identification
 - 2003: response to monkeypox outbreak
- Regulatory agency approval
 - De novo 510K submitted on *Orthopoxvirus* non-variola assay
 - Approved September 2012
 - Discussion initiated with FDA (2002) on *Variola virus* assay
 - Submission on newly validated assays to occur in 2016
- Initiating, technology transfer to other countries
 - Monkeypox
 - Smallpox laboratory diagnostics network (WHO sponsored)

Clinical diagnostic approaches used at the WHO CC at CDC

- Nucleic acid testing
- Viral isolation
- Serologic assays
- Protein based/virus detection
 - In development
 - Commercial assay available
- Only one *Orthopoxvirus* diagnostic assay has achieved regulatory approval
 - LRN *Orthopoxvirus* non-variola real-time PCR assay
 - FDA de novo 510(k) approved September 2012
 - Dependent upon LRN algorithm


Clinical laboratory algorithm development and successes

- Focuses clinical attention to most serious look-a-likes
- Focuses lab attention to most serious contenders
- Helps define/remind what conditions are most frequently confused with possible smallpox
- Frames logic for approaching diagnostics
- Minimizes false positives
- Use of the algorithm in 2002*



LRN

LABORATORY TESTING FOR ACUTE, GENERALIZED VESICULAR OR PUSTULAR RASH ILLNESS IN THE UNITED STATES



Predictive value vs. sensitivity and specificity

CLINICAL DIAGNOSTIC PERFORMANCE RELATED TO DISEASE PREVALENCE

Test parameters

- **Sensitivity:**
 - the ability to identify as positive all those with the disease
- **Specificity:**
 - the ability to identify as negative all those without the disease
- **Predictive value positive (PV+):**
 - the proportion of true positives among those testing positive
- **Predictive value negative (PV-):**
 - the proportion of true negatives among those testing negative

		Disease		total	
		present	absent		
Test result	POS	a=true positives	b=false positives	a+b	PV+ = a/ (a+b)
	NEG	c=false negatives	d=true negatives	c+d	PV- = d/ (c+d)
TOT		a+c	b+d		

$$\text{Sensitivity} = a / (a+c)$$

$$\text{Specificity} = d / (b+d)$$

Pre-event/Post-event

- Test > 90% sensitivity
- Pre-event prevalence of smallpox is zero:
 - If test has 95% specificity, 10 tests done per month, in 6 to 8 sites, every month there will be 3 or 4 false positives
- Post-event prevalence of smallpox is finite:
 - If test has 95% specificity, 1000 tests done per week, every week 50 results will be false positives
 - If test has 99% sensitivity, 1000 tests done per week, 10 results will be false negatives

Test parameters

- Sensitivity and specificity are independent of the prevalence of disease
- Predictive value positive and negative vary with disease prevalence (Bayes' Theorem)
- Implications for smallpox testing "pre-event" and "post-event": use of tests for decision making

Test parameters - examples

- Sensitivity 95%
- Specificity 95%

PREV	PV+	PV-
50%	95%	95%
10%	67.80%	99%
1%	16%	99.95%
0.10%	1.80%	99.99%
0.01%	0.20%	99.99%

Test parameters - examples

- Sensitivity 99%
- Specificity 99%

PREV	PV+	PV-
10.00%	92.80%	99.80%
1%	50%	99.99%
0.10%	9%	100.00%

Example: "Pre-event"

Use more than 1 test to increase PV+

Individual with clinical scenario with fever, followed by centrifugal rash:

	PREV	PV+	PV-
Test 1: sensitivity 99%, specificity 99%	10.00%	92.80%	99.80%
	1%	50%	99.99%
	0.10%	9%	100.00%
	PREV	PV+	PV-
Test 2: sensitivity 95%, specificity 95%	50.00%	95.00%	95.00%
	10%	67%	99.00%
	1.00%	16%	99.50%

Pre event:

prevalence of smallpox is zero

Peri event: prevalence of smallpox is low

- Clinical scenario should be consistent
 - Use febrile rash algorithm; validate algorithm
- Wide availability of other key diagnostic tests especially rapid VZV testing to rule in VZV
- Limit the number of laboratories performing *Variola virus* testing
 - Establish confirmatory testing protocols

**Pre event:
prevalence of smallpox is zero
Peri event: prevalence of smallpox is low**

- Approaches to improving predictive value positive : implications for result use (rule in, and institution of vaccination campaign vs. rule out)
 - Use more than 1 test (different targets)
 - Use tests without common sources of false positives


Post event

- Need to assess what needs for testing will be:
 - One scenario: greatest needs in beginning, and near end of smallpox re-eradication
- May need to test more “low suspicion samples”
 - Sensitivity vs. specificity

Issues relevant to implementation of smallpox diagnostics

- Biosafety:
 - containment;
 - Standard vs. Universal vs. Airborne precautions;
 - vaccination
- Biosecurity
- Reagents
 - WHO, CDC, select agent recommendations on use of *Variola virus* DNA
- Centralized/regionalized testing
 - Transportation of specimens
- QA/QC, Proficiency testing: implementation
- Communication between clinician/epidemiologists/laboratory
 - Clinical history, case patient photos
- Development of a disease confirmatory algorithm
 - Screening tests, confirmatory tests: regional vs. centralized
 - Presumed positive, Confirmed Positive
 - Communication of results, and public health response
 - Role of viral isolation by culture

LRN



Iterative approaches to test validation

NA TEST VALIDATION

Previous validations:

Sensitivity analysis: Serial 100-fold dilutions (1ng/μl to 1 fg/μL) of DNA

Inclusivity panel – *Variola virus* DNA:

- 2 purified viral stocks
- 33 crude viral stocks
- 2 Human scab samples

Specificity analysis: Serial 100-fold dilutions (1ng/μl to 1 fg/μL) of DNA

Exclusivity panel – near neighbor (*Orthopoxvirus*):

- Eurasian: 1 *Ectromelia virus*
- 2 *Monkeypox viruses*
- 2 *Camelpox viruses*
- 1 *Cowpox virus*
- 1 *Taterapox virus*
- 5 *Vaccinia viruses*

North American: 1 *Skunkpox virus*

Exclusivity panel – other rash-causing illnesses:

- 1 *Varicella Zoster virus*
- 1 *Herpes simplex virus* (type 1)
- 1 Rickettsia strain

Exclusivity panel – negatives:

- Myxoma* & tissue culture (2)

Inclusivity Panel		
Species	Strain	Sample
Variola virus	102	Crude
Variola virus	103	Crude
Variola virus	66-39	Crude
Variola virus	7124	Crude
Variola virus	7125	Crude
Variola virus	72-119	Crude
Variola virus	73-175	Crude
Variola virus	77-1605	Crude
Variola virus	Bombay	Crust
Variola virus	Brazil-Garcia	Crude
Variola virus	Congo	Crude
Variola virus	Eth-17	Crude
Variola virus	Harper	Crude
Variola virus	Harvey	Crude
Variola virus	Heidelberg	Crude
Variola virus	Higgins	Crude
Variola virus	Hinton	Crude
Variola virus	Horn	Crude
Variola virus	Horn	Pure
Variola virus	K1629	Crude
Variola virus	Kali Mathu	Crude
Variola virus	Kembula	Crude
Variola virus	Minnesota 124	Crude
Variola virus	MS Lee	Crude
Variola virus	Nepal	Pure
Variola virus	New Dehli	Crude
Variola virus	Nigeria Kuciano	Crust
Variola virus	Nur Islam	Crude
Variola virus	Rumbec	Crude
Variola virus	Shahzamon	Crude
Variola virus	Solaiman	Crude
Variola virus	Stillwell	Crude
Variola virus	V68-59	Crude
Variola virus	V70-222	Crude
Variola virus	V70-228	Crude
Variola virus	Variolator-4	Crude
Variola virus	Yamada	Crude

Complementary set:


Exclusivity Panel - near neighbor (<i>Orthopoxviruses</i>)	
Species	Strain
<i>Ectromelia virus</i>	Moscow
<i>Monkeypox virus</i>	79-0266
<i>Monkeypox virus</i>	79-0005
<i>Camelpoxvirus</i>	LLC
<i>Camelpoxvirus</i>	V78-1-903
<i>Cowpoxvirus</i>	Brighton
<i>Taterapoxvirus</i>	(Gerbilpox)
<i>Vaccinia virus</i>	Lister
<i>Vaccinia virus</i>	VTH
<i>Vaccinia virus</i>	Wyeth
<i>Vaccinia virus</i>	WYH pGS62-9-v1-1-1
<i>Vaccinia virus</i>	Rabbitpoxvirus
<i>Skunkpoxvirus</i>	

Exclusivity Panel - other rash-causing illnesses	
Species	Strain
<i>Varicella Zoster Virus</i>	Webster
<i>Herpes Simplex Virus-1</i>	HFEM
<i>Rickettsia</i>	conorii

Exclusivity Panel - other negatives	
Species	Strain
<i>Myxoma</i>	
Human tissue culture cells	Sup-T
Monkey kidney tissue culture cells	BSC-40

Variola virus : diagnostic creation and validation

- ❑ Validated diagnostic real-time PCR assays
 - Target multiple regions of the genome
 - Target sequences specific to *Variola virus*
- ❑ *Cowpox virus* sequences acquired after validation
 - Exhibit extensive phylogenetic diversity
 - Contain certain regions previously thought to be specific to *Variola virus*
 - Now assays predicted to cross-react with *Cowpox virus*



D. Carroll, et. al

Variola virus signature (eroded specificity): assay cross-reacts with Cowpox virus

Exclusivity panel Species	Strain Name	Assay 1 ct for 5 ng DNA	Assay 2 ct for 5 ng DNA
<i>Vaccinia</i>	Copenhagen	Negative	Negative
<i>Vaccinia</i>	WR	Negative	Negative
<i>Vaccinia</i>	ACAM 2000	Negative	Negative
<i>Vaccinia</i>	BRZ SERRO	Negative	Negative
<i>Cowpox</i>	CPXV-NOR1995-MAN	Negative	Negative
<i>Cowpox</i>	CPXV GER1980-EP4	19	Negative
<i>Cowpox</i>	CPXV GER1991-3	18	Negative
<i>Cowpox</i>	CPXV GER1998_2	17	Negative
<i>Cowpox</i>	CPXV FIN 2000		Negative
<i>Ectromelia</i>	ECTV Moscow	Negative	Negative
<i>Monkeypox</i>	MPXV RCG 2003 358	Negative	Negative
<i>Monkeypox</i>	MPXV USA 2003 044	Negative	Negative
<i>Raccoonpox</i>	RACV V71-I-84	Negative	Negative
<i>Skunkpox</i>	SKPV 1991	Negative	Negative
<i>Volepox</i>	VPXV 2004-CA-007	Negative	Negative
<i>Camelpox</i>	CMLV-78-I-2379	17	Negative
<i>Taterapox (gerbilpox)</i>	TATV-71-I-016	16	Negative
<i>Parapoxvirus Orf</i>	Vaccine for sheep	Negative	Negative

What we have learned about *Variola virus* diagnostic assay development/validation

- ❑ Bioinformatic analysis should lead design of validation panels
 - Inclusivity panel include all *Variola virus* strains with differences in assay target region
 - Exclusivity panel (near neighbor *Orthopoxvirus*) contain viruses with assay target regions most similar to *Variola virus*
 - Exceedingly difficult to construct uniform panels for all assays due to high similarity between *Orthopoxviruses*
- ❑ Simultaneous identification of multiple *Variola virus* signatures will increase confidence in initial identification/verification of the pathogen with real-time PCR

Current recommendations: tailor panel to assay based on bioinformatics

Sensitivity analysis: Two low dilutions (100 and 10 fg/μL) of DNA
Inclusivity panel – *Variola virus*: ≥ 2 purified viral stocks
(at least one from each primary clade)
(encompass differences in target region)

Specificity analysis: One high dilution (100 pg/μl) of nucleic acid
Exclusivity panel – near neighbor (*Orthopoxvirus*):
Eurasian: 1 *Ectromelia virus*
2 *Monkeypox viruses* (one from each clade)
1 *Camelpox virus*
5 *Cowpox viruses* (one from each predicted clade)
1 *Taterapox virus*
4 *Vaccinia viruses* (one from each clade)
North American: 1 *Raccoonpox virus*
1 *Skunkpox virus*
1 *Volepox virus*

Current recommendations: tailor panel to assay based on bioinformatics

Specificity analysis (cont.): One high dilution (100 pg/μl) of DNA

Exclusivity panel (cont.) – other rash-causing illnesses:
2 *Varicella Zoster viruses* (J clade and E1 clade)
2 *Herpes simplex viruses* (type 1 and 2)
2 Rickettsia strains

Chordopoxviruses:
1 *Parapoxvirus (Orf)*

Negatives:
13 bacteria strains (skin flora)
water

Current Recommendations:

Exclusivity panels
100 pg/ μ l

Exclusivity panel - near neighbors (Orthopoxviruses)	
Species	Strain Name
<i>Ectromelia</i>	ECTV Moscow
<i>Monkeypox</i>	MPXV RCG 2003 358
<i>Monkeypox</i>	MPXV USA 2003 044
<i>Camelpox</i>	CMLV-78-I-2379
<i>Cowpox</i>	CPXV-NOR1995-MAN
<i>Cowpox</i>	CPXV GER1980-EP4
<i>Cowpox</i>	CPXV GER1991-3
<i>Cowpox</i>	CPXV FIN-2000-MAN
<i>Cowpox</i>	CPXV GER1998 2
<i>Taterapox (gerbilpox)</i>	TATV-71-I-016
<i>Vaccinia</i>	Copenhagen
<i>Vaccinia</i>	WR
<i>Vaccinia</i>	ACAM 2000
<i>Vaccinia</i>	BRZ SERRO
<i>Raccoonpox</i>	RACV V71-I-84
<i>Skunkpox</i>	SKPV 1991
<i>Volepox</i>	VPXV 2004-CA-007

Exclusivity panel - other rash-causing illnesses

Species	Strain Name
<i>Varicella-zoster virus</i>	pOKA (J clade)
<i>Varicella-zoster virus</i>	Webster (E1 clade)
<i>Herpes simplex virus type 1</i>	F
<i>Herpes simplex virus type 2</i>	G
<i>Rickettsia conorii</i>	CDC
<i>Rickettsia akari</i>	CDC
<i>Parapoxvirus Orf</i>	Vaccine for sheep

Exclusivity panel - Negatives

Species	ID number
<i>Enterococcus faecalis</i>	ATCC 29212
<i>Eschericia coli</i>	ATCC 25922
<i>Klebsiella pneumoniae</i>	ATCC 33495
<i>Peptostreptococcus anaerobius</i>	ATCC 27337
<i>Propionibacterium acnes</i>	ATCC 6919
<i>Pseudomonas aeruginosa</i>	ATCC 27853
<i>Staphylococcus aureus</i> (strain 1)	ATCC 12600
<i>Staphylococcus aureus</i> (strain 2)	ATCC 25923
<i>Staphylococcus epidermidis</i> (strain 1)	ATCC 49134
<i>Staphylococcus epidermidis</i> (strain 2)	ATCC 12228
<i>Staphylococcus epidermidis</i> (strain 3)	ATCC 14990
<i>Streptococcus gallylyticus</i>	ATCC 49147
<i>Streptococcus pyogenes</i>	ATCC 49117
Water	

**THINKING ABOUT ENVIRONMENTAL
DETECTION**

Considerations in addition to those of clinical diagnostics....

- ❑ **How to verify authentic agent?**
 - Sufficient vs. necessary
 - Culture?
 - Nucleic acid tests
 - How much of the genome?
- ❑ **How to verify infectious risk?**
 - Absent a clinically ill human, what is sufficient, what is necessary?
- ❑ **How and where to evaluate validate**
 - WHO considerations what is sufficient, necessary to have a public safety/health actionable assay result;
 - What should that action be?

Mission of the SPADA Variola Working Group (2014)

- The Variola Working Group of the Stakeholder Panel on Agent Detection Assays (SPADA) was tasked to develop voluntary consensus standards required for evaluation of tools that detect Variola from aerosol collection samples. ...
The standards will :
 - Support test and evaluation of Variola-detection tools for the BioWatch program
 - Provide guidance to industry and other capability developers for development of future detection tools that BioWatch may solicit

It is expected that any detection result from a tool validated against the SPADA Variola standards will be confirmed by the Poxvirus Laboratory at the Centers for Disease Control and Prevention.

Variola Working Group SMPR: tailor panel to assay based on bioinformatics

Sensitivity analysis:

- AMDL = 50,000 copies/mL target region of *Variola virus*
 - ≥ 500 bp must receive permission from WHO
 - insertion into another *Orthopoxvirus* is prohibited
- Establish Probability of Detection at AMDL w/in collection buffer (≥ 95%)
- Establish Probability of Detection at AMDL w/in environmental matrix (≥ 95%)
- Inclusivity panel – *Variola virus*: ≥ 2 strain target regions
 - at least one from each primary clade
 - encompass differences in target region
 - Based on bioinformatic analysis
- Ensure all inclusivity strains are detected at 2X AMDL in collection buffer
- Ensure all inclusivity strains are detected at 2X AMDL in environmental matrix
- Ensure target is detected at 2X AMDL w/in pool of environmental panel organisms (pools of up to 10 organisms at 10X AMDL for each)

Variola Working Group SMPR: tailor panel to assay based on bioinformatics

Specificity analysis:

- Exclusivity panel – near neighbor (*Orthopoxvirus*):
 - All poxvirus strains listed in the table (one from each major clade)
 - See AOAC Website for the most updated list

Species	Strain	Commercial availability
<i>Vaccinia</i>	Elsree	ATCC VR-1549
<i>Cowpox</i>	Brighton	ATCC VR-302
<i>Ectromelia</i>	Moscow	ATCC VR-1374
<i>Monkeypox</i>	V79-I-005	BEI NR-2324
<i>Monkeypox</i>	USA-2003	BEI NR-2500
<i>Raccoonpox</i>	Herman	ATCC VR-838
<i>Skunkpox</i>		ATCC
<i>Volepox</i>		ATCC
<i>Camelpox</i>		BEI
<i>Taterapox</i>		BEI
<i>Parapoxvirus Orf</i>	Vaccine	Colorado Serum Company

- Any additional strains with greater similarity to the assay's target region(s) than the strains listed above in the table
 - Based on bioinformatic analysis
- Ensure all exclusivity strains are NOT detected at 10X AMDL in collection buffer

Variola Working Group SMPR: tailor panel to assay based on bioinformatics

Specificity analysis (cont.):

- Environmental aerosol matrix samples:
 - Method developers should obtain environmental matrix samples that are representative/consistent with the collection method to be used
 - Considerations include:
 - Collection medium
 - Duration of collection
 - Diversity of geographical areas to be sampled
 - Climatic/environmental conditions
 - Seasonal changes
 - Ensure sufficient replicates to represent environmental condition
 - Ensure aerosol matrix samples do NOT cross-react
- Environmental panel organisms:
 - Organisms can be pooled (up to 10 per pool)
 - Method developer must justify exclusion of specific panel organisms
 - Ensure all organisms are NOT detected at 10X AMDL
 - If unexpected result, each individual organisms from failed pool must be tested individually at 10X AMDL

Variola Working Group SMPR: tailor panel to assay based on bioinformatics

Bioinformatic analysis:

- *In silico* screening on signature sequences
 - Suggestive of potential performance issues
 - Guide necessary additions to wet lab screening panels
- Potential tools for *in silico* screening:
 - Simulate_PCR
 - NCBI tools
 - FastPCR
- Method developer submission should include:
 - Description of sequence databases used in the *in silico* analysis
 - Description of tool used for bioinformatics evaluation
 - Data confirming selected tool performance based on wet-lab testing
 - Can be generated retrospectively using published assays
 - List of additional strains to be added to inclusivity or exclusivity panels

Acknowledgements

Poxvirus and Rabies Branch members

1999-present

"TNTC" CDC and external partners

For more information please contact Centers for Disease Control and Prevention

1600 Clifton Road NE, Atlanta, GA 30333
 Telephone, 1-800-CDC-INFO (232-4636)/TTY: 1-888-232-6348
 E-mail: cdcinfo@cdc.gov Web: www.cdc.gov

The findings and conclusions in this report are those of the authors and do not necessarily represent the official position of the Centers for Disease Control and Prevention.

National Center for Emerging and Zoonotic Infectious Diseases
 Division of High Consequence Pathogens and Pathology



Environmental Variola screening test positives (26)

>808,000 samples tested

confirmatory testing: none confirmed positive

Test Date	Assay	Sample ID	Parameter	Result	CT Value
2/10/2008 08:09:00	PM CDC Screen Panel	P066079	VRL3	reactive	40.54
2/10/2008 08:07:00	PM CDC Screen Panel	P066060	VRL3	reactive	40.54
2/10/2008 10:18:00	PM Verification panel for Variola virus	P066060	VRL1	reactive	27.89
2/10/2008 08:11:00	PM CDC Screen Panel	P065784	VRL3	reactive	40.54
2008					
5/6/2009 02:43:01	PM CDC Screen Panel	D50609-654	VRL3	reactive	28.93
5/6/2009 02:43:00	PM CDC Screen Panel	D50609-192	VRL3	reactive	28.93
5/6/2009 02:43:01	PM CDC Screen Panel	D50609-666	VRL3	reactive	28.93
11/23/2009 01:16:00	PM CDC Screen Panel	P069407	VRL3	reactive	41.69
11/23/2009 01:16:00	PM CDC Screen Panel	P069411	VRL3	reactive	41.69
12/17/2009 03:34:00	PM CDC Screen Panel	P070763	VRL3	reactive	43.28
12/17/2009 03:34:00	PM CDC Screen Panel	P070670	VRL3	reactive	43.28
10/9/2009 03:21:00	PM CDC Screen Panel	P133565	VRL3	reactive	20.66
2009					
4/17/2010 08:37:00	PM CDC Screen Panel	P298637	VRL3	reactive	28.91
4/17/2010 08:37:00	PM CDC Screen Panel	P298640	VRL3	reactive	28.36
4/17/2010 08:37:00	PM CDC Screen Panel	P298645	VRL3	reactive	28.03
4/26/2010 02:13:00	PM CDC Screen Panel	P222219	VRL3	reactive	42.28
4/3/2010 03:40:00	PM CDC Screen Panel	P113811	VRL3	reactive	36.92
4/3/2010 03:40:00	PM CDC Screen Panel	P113819	VRL3	reactive	36.92
2/23/2010 01:03:00	PM CDC Screen Panel	D549214	VRL3	reactive	44.07
3/29/2010 09:48:00	AM CDC Screen Panel	P554250	VRL3	reactive	41.02
3/29/2010 09:48:00	AM CDC Screen Panel	P554251	VRL3	reactive	41.02
3/29/2010 09:48:00	AM CDC Screen Panel	P554252	VRL3	reactive	41.02
3/17/2010 02:50:00	PM CDC Screen Panel	P076217	VRL3	reactive	26.53
3/17/2010 02:50:00	PM CDC Screen Panel	P076225	VRL3	reactive	26.53
8/5/2010 07:33:00	PM CDC Screen Panel	P409223	VRL3	reactive	42.87
2010					
12/29/2011 07:47:00	PM CRP Screen Panel	P307499	POX2	reactive	32.74
12/29/2011 07:43:00	PM CRP Screen Panel	P307493	POX2	reactive	32.74
7/21/2011 07:39:00	PM CDC Screen Panel	P066838	VRL3	reactive	44.01
1/7/2011 02:20:00	PM CDC Screen Panel	P194059	VRL3	reactive	44.01
2011					
					4
					29

AOAC Acronyms and abbreviations

AMDL	acceptable minimum detection level
AOAC	AOAC INTERNATIONAL (AOAC formerly stood for <i>Association of Official Analytical Chemists</i> , but long-name no longer used)
CSO	chief scientific officer
ERP	expert review panel
ISO	International Organization for Standardization
LOD	limit of detection
LPOD	laboratory probability of detection
NGO	non-governmental organization
OMA	<i>Official Methods of Analysis</i> , frequently pronounced like “o maa”
POD	probability of detection
SPADA	Stakeholder Panel on Agent Detection Assays
SMPR	Standard Method Performance Requirements, frequently pronounced as in “smipper”.



March 16, 2016

STAKEHOLDER PANEL ON AGENT DETECTION ASSAYS

RESOURCES

SPDS

Key Staff Contacts:

Name	Role	Email	Telephone
Scott Coates	AOAC Chief Scientific Officer	scoates@aoac.org	301.924.7077 x 137
Christopher Dent	Standards Development Coordinator	cdent@aoac.org	301.924.7077 x 119
Krystyna McIver	Executive, Scientific Business Development (SPADA Project Executive)	kmciver@aoac.org	301.924.7077 x 111
Deborah McKenzie	Sr. Director, Standards Development and Method Approval Processes	dmckenzie@aoac.org	301.924.7077 x 157

AOAC Website: <http://www.aoac.org>

SPADA Website: <http://bit.ly/1Hmf6ba>

SPADA Working Group Sign Up: <https://form.jotform.com/53514168225150>



SPADA

Role	Name	Organization
Chair	Linda Beck	CBR Defense Concepts And Experimentation Branch, Naval Surface Warfare Center
Chair	Matthew Davenport	Department Of Homeland Security
SPADA Member	James Agin	Q Laboratories, Inc.
SPADA Member	Terrance Allen	Pentagon Force Protection Agency
SPADA Member	Amy Altman	Luminex Corporation
SPADA Member	Douglas Anders	Federal Bureau Of Investigation (FBI)
SPADA Member	Jessica Appler	HHS BARDA
SPADA Member	Jennifer Arce	PNNL
SPADA Member	Thomas Archibald	HazTech Systems, Inc
SPADA Member	Charles Asowata	Executive Office For Chemical And Biological Defense
SPADA Member	Les Baillie	School Of Pharmacy And Pharmaceutical Sciences, Cardiff University
SPADA Member	Ed Bailor	IAB
SPADA Member	Jeff Ballin	ECBC
SPADA Member	Timothy Bauer	Naval Surface Warfare Center Dahlgren
SPADA Member	Maureen Beanan	National Institutes Of Health
SPADA Member	Brian Bennett	West Desert Test Center, CAPAT, Dugway Proving Ground
SPADA Member	Thomas Blank	NBACC
SPADA Member	Steven Blanke	University of Illinois
SPADA Member	Jerold Blutman	DTRA
SPADA Member	Larry Blyn	Ibis Biosciences
SPADA Member	Donna Boston	HHS, ASPR/BARDA
SPADA Member	Julie Boylan	Defense Threat Reduction Agency
SPADA Member	Carrie Brennan	Austin Peay State University
SPADA Member	Paul Brett	University Of South Alabama
SPADA Member	Cindy Bruckner-Lea	Pacific Northwest National Lab
SPADA Member	Robert Bull	Department Of Homeland Security
SPADA Member	Mary Burtnick	University Of South Alabama
SPADA Member	Don Bushner	JS J8, JRO-CBRND
SPADA Member	Ryan Cahall	Censeo Insight
SPADA Member	Andrew Cannon	USF Center For Biological Defense
SPADA Member	Amanda Clark	Naval Surface Warfare Center Dahlgren Virginia
SPADA Member	Kenneth Cole	NIST
SPADA Member	Bart Currie	Tropical And Emerging Infectious Diseases Division, Menzies School Of Health Res

Don't see your name on this list? Let us know at cdent@aoac.org to ensure you receive SPADA emails!

Role	Name	Organization
SPADA Member	Kenneth Damer	Northrop Grumman Electronic Systems
SPADA Member	Inger Damon	Centers For Disease Control And Prevention
SPADA Member	David Dance	Health Protection Agency (South West)
SPADA Member	David DeShazer	USAMRIID
SPADA Member	Christina Egan	NYSDOH
SPADA Member	Phillip Elzer	LSU AgCenter
SPADA Member	Peter Emanuel	BioScience
SPADA Member	Thomas Ficht	TAMU
SPADA Member	Brandan Fisher	US Army Test And Evaluation Command
SPADA Member	Steven Fisher	The Office Of Justice Programs (OJP)
SPADA Member	Bill Folkerts	ATCC
SPADA Member	Mats Forsman	FOI Sweden
SPADA Member	Edward Gabriel	DHS - HHS
SPADA Member	Ken Gage	CDC
SPADA Member	Mike Gavin	Emergency Manager City Of Fort Collins
SPADA Member	Joan Gebhardt	Naval Medical Research Center
SPADA Member	Jay Gee	CDC
SPADA Member	Jennifer Gibbons	ECBC / Excet
SPADA Member	Michael Gillenwater	DHS - FEMA, Preparedness Grants Division
SPADA Member	Arthur Goff	CIV USA MEDCOM USAMRIID
SPADA Member	Bruce Goodwin	DoD CRP
SPADA Member	Robert Griffin	DHS S&T First Responders Group
SPADA Member	Ted Hadfield	HADECO, LLC
SPADA Member	Martha Hale	US ARMY MEDCOM USAMRIID
SPADA Member	Andrew Hebbeler	Office Of Science And Technology Policy
SPADA Member	Matthew Hickman	Department Of Homeland Security
SPADA Member	Steven Hinrichs	University Of Nebraska Medical Center
SPADA Member	Anthony Hitchins	FDA - CFSAN (Retired)
SPADA Member	Peyton Hobson	FDA HHS
SPADA Member	David Hodge	Department Of Homeland Security
SPADA Member	Alex Hoffmaster	Centers For Disease Control And Prevention
SPADA Member	Jeffery Hogan	ATEC, Life Sciences Division, Dugway Proving Ground
SPADA Member	Harvey Holmes	Centers for Disease Control

Don't see your name on this list? Let us know at cdent@aoac.org to ensure you receive SPADA emails!

Role	Name	Organization
SPADA Member	Kia Hopkins	ECBC
SPADA Member	Rosemary Humes	HHS, ASPR/BARDA
SPADA Member	Duane Hunt	Baltimore City Environmental Services
SPADA Member	Aaron Hyre	JBTDs/NBCCA
SPADA Member	Sofi Ibrahim	USAMRIID
SPADA Member	Paula Imbro	The Tauri Group
SPADA Member	Robert Ingram	FDNY
SPADA Member	Paul Jackson	Lawrence Livermore National Lab (Retired)
SPADA Member	Crystal Jaing	LLNL
SPADA Member	Malcolm Johns	DHS
SPADA Member	Frederick Johnson	DA DCS G-3/5/7
SPADA Member	Ronald Johnson	BioMérieux, Inc.
SPADA Member	Franca Jones	White House Office of Science and Technology Policy
SPADA Member	Brian Kamoie	DHS - FEMA - Grants Program Directorate
SPADA Member	Cecilia Kato	CDC
SPADA Member	Alexander Kayatani	Pentagon Force Protection Agency
SPADA Member	Paul Keim	Northern Arizona University
SPADA Member	Liz Kerrigan	ATCC
SPADA Member	Saleem Khan	University Of Pittsburgh School Of Medicine
SPADA Member	Katalin Kiss	ATCC
SPADA Member	Kristin Korte	ICx Technologies
SPADA Member	Matt Kramer	Qiagen
SPADA Member	Tom Labombarda	Aventura Police Department
SPADA Member	Markus Lacorn	R-Biopharm AG
SPADA Member	David Ladd	The Commonwealth Of Massachusetts Department Of Fire Services
SPADA Member	John Lednický	University Of Florida
SPADA Member	Matthew Lesho	Luminex
SPADA Member	Direk Limmathurotsakul	Mahidol-Oxford Tropical Medicine Research Unit, Thailand
SPADA Member	Nancy Lin	NIST
SPADA Member	Luther Lindler	DHS
SPADA Member	Angelo Madonna	Dugway Proving Ground - Life Sciences Division
SPADA Member	Chris Mangal	Association Of Public Health Laboratories
SPADA Member	Laura Maple	NSWC

Don't see your name on this list? Let us know at cdent@aoac.org to ensure you receive SPADA emails!

Role	Name	Organization
SPADA Member	James Marks	UCSF
SPADA Member	Jim Marks	UCSF
SPADA Member	Bryon Marsh	Georgia National Guard
SPADA Member	Robert Massung	Centers For Disease Control And Prevention
SPADA Member	J. Clay McGuyer	National Guard Bureau
SPADA Member	Marian McKee	BioReliance Corporation
SPADA Member	Michael McLaughlin	US FDA - ORA/ORS
SPADA Member	Toby Merlin	Centers For Disease Control And Prevention
SPADA Member	Richard Meyer	
SPADA Member	Timothy Minogue	DoD, USAMRIID
SPADA Member	Jayne Morrow	NSTC
SPADA Member	Stephen Morse	Centers For Disease Control And Prevention(Retired)
SPADA Member	Timothy Moshier	Acumen Detection LLC
SPADA Member	Pejman Naraghi-Arani	InSilixa Corp.
SPADA Member	Dallas New	
SPADA Member	Ann Nguyen	BARDA
SPADA Member	Christopher Niblick	JPM NBC CA, PD CCAT&TI
SPADA Member	William Nierman	J. Craig Venter Institute
SPADA Member	Mikeljon Nikolich	Walter Reed Army Institute Of Research
SPADA Member	Sean O'Brien	DoD DUSA T&E
SPADA Member	Tom O'Brien	Tetracore, Inc.
SPADA Member	Catherine O'Connell	Life Technologies
SPADA Member	William O'Neill	US Postal Service
SPADA Member	Steven Olsen	USDA ARS
SPADA Member	Ted Olsen	PathSensors, Inc.
SPADA Member	Victoria Olson	Centers For Disease Control And Prevention
SPADA Member	Kate Ong	JPEO-CBD
SPADA Member	Palmer Orlandi	FDA Office of Foods and Veterinary Medicine
SPADA Member	Eileen Ostlund	USDA, APHIS, Veterinary Services
SPADA Member	Rich Ozanich	Pacific Northwest National Laboratory
SPADA Member	Traci Pals	DTRA
SPADA Member	Joseph Perrone	SRI International Bioscience Division
SPADA Member	Robert Perry	University Of Kentucky

Don't see your name on this list? Let us know at cident@aoac.org to ensure you receive SPADA emails!

Role	Name	Organization
SPADA Member	Jeannine Petersen	Centers For Disease Control And Prevention (CDC)
SPADA Member	Denise Pettit	N. C. Department Of Health And Human Services
SPADA Member	Sally Phillips	DHS OHA
SPADA Member	Mark Poli	USAMRIID
SPADA Member	Mark Poli	US Army Medical Research Institute Of Infectious Diseases
SPADA Member	Ann Powers	CDC
SPADA Member	Lyle Probst	Microfluidic Systems, Inc.
SPADA Member	Richard Prouty	JRO-CBRND
SPADA Member	Amy Pullman	Association Of Public Health Laboratories
SPADA Member	Beena Puri	FDA Division Of Microbiology
SPADA Member	Amy Rasley	Lawrence Livermore National Laboratory
SPADA Member	Jon Rayner	Southern Research Institute
SPADA Member	Roberto Rebeil	ECBC
SPADA Member	Christopher Renner	JRO-CBRND
SPADA Member	Michael Retford	JBTDJ JPM NBCCA
SPADA Member	Frank Roberto	Idaho National Laboratory
SPADA Member	Frank Roberto	Idaho National Laboratory
SPADA Member	Jason Roos	JPEO-CBD
SPADA Member	David Rozak	USAMRIID
SPADA Member	James Samuel	Texas A&M
SPADA Member	Stefan Saravia	Minnesota Department Of Health Public Health Laboratory Division
SPADA Member	Frank Schaefer	US EPA (ret)
SPADA Member	Mark Scheckelhoff	DHS/OHA
SPADA Member	Herbert Schweizer	University of Florida
SPADA Member	Sanjiv Shah	US EPA
SPADA Member	Shashi Sharma	FDA - CFSAN
SPADA Member	Deborah Shuping	DoD, DUSA-TE
SPADA Member	Russell Sillmon	CTTSO
SPADA Member	Tom Slezak	Lawrence Livermore National Lab
SPADA Member	Darci Smith	Southern Research Institute
SPADA Member	Theresa Smith	USAMRIID
SPADA Member	Sandra Smole	Massachusetts Department Of Public Health
SPADA Member	Shanmuga Sozhamannan	DoD ECBC

Don't see your name on this list? Let us know at cdent@aoac.org to ensure you receive SPADA emails!

Role	Name	Organization
SPADA Member	Darryl Sullivan	Covance Laboratories
SPADA Member	Maureen Sullivan	Minnesota Department Of Health
SPADA Member	Mary Beth Tabacco	Smiths Detection
SPADA Member	Sandra Tallent	FDA - ORS/DM
SPADA Member	Christina Thompson	Thompson Biosafety, LLC
SPADA Member	Maureen Thornton	MRI Global
SPADA Member	Rebekah Tiller	CDC
SPADA Member	Patrick Treado	ChemImage Corp
SPADA Member	David Trudil	New Horizons Diagnostics Corporation
SPADA Member	APICHAJ TUANYOK	University of Florida
SPADA Member	Venkat Venkateswaran	Omni Array Biotechnology
SPADA Member	Elizabeth Vitalis	Lawrence Livermore National Laboratory
SPADA Member	David Wagner	Northern Arizona University
SPADA Member	Jennifer Walker	Tetracore Inc.
SPADA Member	Rodney Wallace	BARDA
SPADA Member	Mike Walter	Office Of Health AffairsDHS
SPADA Member	Richard Warren	BARDA/Contractor Tunnell Government Services
SPADA Member	David Watson	DTRA
SPADA Member	Scott Weaver	University Of Texas Medical Branch
SPADA Member	Clyde Webster	DoD DUSA T&E
SPADA Member	Susan Welkos	USAMRIID
SPADA Member	James Whelan	Alexeter Technologies
SPADA Member	Christian Whitchurch	DTRA
SPADA Member	Patrick Williams	Evogen
SPADA Member	Reinhardt Witzemberger	R-Biopharm AG
SPADA Member	Donald Woodbury	DHS CDB S&T
SPADA Member	Patricia Worsham	USAMRIID
SPADA Member	Emily Yost	ATEC, Life Sciences Division, Dugway Proving Ground
SPADA Member	Edward Young	VA Medical Center/Baylor College Of Medicine
OMB Advisor	Douglas Abbott	USDA - FSIS/OPHS/BD/MOSPB
AOAC Staff	Scott Coates	AOAC INTERNATIONAL
AOAC Staff	Christopher Dent	AOAC INTERNATIONAL
AOAC Staff	Krystyna McIver	AOAC INTERNATIONAL

Don't see your name on this list? Let us know at cdent@aoac.org to ensure you receive SPADA emails!

Role	Name	Organization
AOAC Staff	Deborah McKenzie	AOAC INTERNATIONAL
AOAC Staff	Tien Milor	AOAC INTERNATIONAL
OMB Advisor	Tom Phillips	MD Department Of Agriculture

Appendix W

POLICY AND PROCEDURES ON VOLUNTEER CONFLICT OF INTEREST

Statement of Policy

While it is not the intention of AOAC INTERNATIONAL (AOAC) to restrict the personal, professional, or proprietary activities of AOAC members nor to preclude or restrict participation in Association affairs solely by reason of such activities, it is the sense of AOAC that conflicts of interest or even the appearance of conflicts of interest on the part of AOAC volunteers should be avoided. Where this is not possible or practical under the circumstances, there shall be written disclosure by the volunteers of actual or potential conflicts of interest in order to ensure the credibility and integrity of AOAC. Such written disclosure shall be made to any individual or group within the Association which is reviewing a recommendation which the volunteer had a part in formulating and in which the volunteer has a material interest causing an actual or potential conflict of interest.

AOAC requires disclosure of actual or potential conflicts of interest as a condition of active participation in the business of the Association. The burden of disclosure of conflicts of interest or the appearance of conflicts of interest falls upon the volunteer.

A disclosed conflict of interest will not in itself bar an AOAC member from participation in Association activities, but a three-fourths majority of the AOAC group reviewing the issue presenting the conflict must concur by secret ballot that the volunteer's continued participation is necessary and will not unreasonably jeopardize the integrity of the decision-making process.

Employees of AOAC are governed by the provision of the AOAC policy on conflict of interest by staff. If that policy is in disagreement with or mute on matters covered by this policy, the provisions of this policy shall prevail and apply to staff as well.

Illustrations of Conflicts of Interest

1. A volunteer who is serving as a committee member or referee engaged in the evaluation of a method or device; who is also an employee of or receiving a fee from the firm which is manufacturing or distributing the method or device or is an employee of or receiving a fee from a competing firm.
2. A volunteer who is requested to evaluate a proposed method or a related collaborative study in which data are presented that appear detrimental (or favorable) to a product distributed or a position supported by the volunteer's employer.
3. A referee who is conducting a study and evaluating the results of an instrument, a kit, or a piece of equipment which will be provided gratis by the manufacturer or distributor to one or more of the participating laboratories, including his or her own laboratory, at the conclusion of the study.
4. Sponsorship of a collaborative study by an interest (which may include the referee) which stands to profit from the results; such sponsorship usually involving the privilege granted by the investigator to permit the sponsor to review and comment upon the results prior to AOAC evaluation.
5. A volunteer asked to review a manuscript submitted for publication when the manuscript contains information which is critical of a proprietary or other interest of the reviewer.

The foregoing are intended as illustrative and should not be interpreted to be all-inclusive examples of conflicts of interest AOAC volunteers may find themselves involved in.

Do's and Don't's

Do avoid the appearance as well as the fact of a conflict of interest.

Do make written disclosure of any material interest which may constitute a conflict of interest or the appearance of a conflict of interest.

Do not accept payment or gifts for services rendered as a volunteer of the Association without disclosing such payment or gifts.

Do not vote on any issue before an AOAC decision-making body where you have the appearance of or an actual conflict of interest regarding the recommendation or decision before that body.

Do not participate in an AOAC decision-making body without written disclosure of actual or potential conflicts of interest in the issues before that body.

Do not accept a position of responsibility as an AOAC volunteer, without disclosure, where the discharge of the accepted responsibility will be or may appear to be influenced by proprietary or other conflicting interests.

Procedures

Each volunteer elected or appointed to an AOAC position of responsibility shall be sent, at the time of election or appointment, a copy of this policy and shall be advised of the requirement to adhere to the provisions herein as a condition for active participation in the business of the Association. Each volunteer, at the time of his or her election or appointment, shall indicate, in writing, on a form provided for this purpose by AOAC, that he or she has read and accepts this policy.

Each year, at the spring meeting of the AOAC Board of Directors, the Executive Director shall submit a report certifying the requirements of this policy have been met; including the names and positions of any elected or appointed volunteers who have not at that time indicated in writing that they have accepted the policy.

Anyone with knowledge of specific instances in which the provisions of this policy have not been complied with shall report these instances to the Board of Directors, via the Office of the Executive Director, as soon as discovered.

* * * * *

Adopted: March 2, 1989

Revised: March 28, 1990

Revised: October 1996

Reviewed by outside counsel March 2000 (Fran Dwornik) and found to be current and relevant

Appendix U

ANTITRUST POLICY STATEMENT AND GUIDELINES

Introduction

It is the policy of AOAC INTERNATIONAL (AOAC) and its members to comply strictly with all laws applicable to AOAC activities. Because AOAC activities frequently involve cooperative undertakings and meetings where competitors may be present, it is important to emphasize the on-going commitment of our members and the Association to full compliance with national and other antitrust laws. This statement is a reminder of that commitment and should be used as a general guide for AOAC and related individual activities and meetings.

Responsibility for Antitrust Compliance

The Association's structure is fashioned and its programs are carried out in conformance with antitrust standards. However, an equal responsibility for antitrust compliance -- which includes avoidance of even an appearance of improper activity -- belongs to the individual. Even the appearance of improper activity must be avoided because the courts have taken the position that actual proof of misconduct is not required under the law. All that is required is whether misconduct can be inferred from the individual's activities.

Employers and AOAC depend on individual good judgment to avoid all discussions and activities which may involve improper subject matter and improper procedures. AOAC staff members work conscientiously to avoid subject matter or discussion which may have unintended implications, and counsel for the Association can provide guidance with regard to these matters. It is important for the individual to realize, however, that the competitive significance of a particular conduct or communication probably is evident only to the individual who is directly involved in such matters.

Antitrust Guidelines

In general, the U.S. antitrust laws seek to preserve a free, competitive economy and trade in the United States and in commerce with foreign countries. Laws in other countries have similar objectives. Competitors (including individuals) may not restrain competition among themselves with reference to the price, quality, or distribution of their products, and they may not act in concert to restrict the competitive capabilities or opportunities of competitors, suppliers, or customers.

Although the Justice Department and Federal Trade Commission generally enforce the U.S. antitrust laws, private parties can bring their own lawsuits. Penalties for violating the U.S. and other antitrust laws are severe: corporations are subject to heavy fines and injunctive decrees, and may have to pay substantial damage judgments to injured competitors, suppliers, or customers. Individuals are subject to criminal prosecution, and will be punished by fines and imprisonment. Under current U.S. federal sentencing guidelines, individuals found guilty of bid rigging, price fixing, or market allocation must be sent to jail for at least 4 to 10 months and must pay substantial minimum fines.

Since the individual has an important responsibility in ensuring antitrust compliance in AOAC activities, everyone should read and heed the following guidelines.

1. Don't make any effort to bring about or prevent the standardization of any method or product for the purpose or intent of preventing the manufacture or sale of any method or product not conforming to a specified standard
2. Don't discuss with competitors your own or the competitors' prices, or anything that might

- affect prices such as costs, discounts, terms of sale, distribution, volume of production, profit margins, territories, or customers.
3. Don't make announcements or statements at AOAC functions, outside leased exhibit space, about your own prices or those of competitors.
 4. Don't disclose to others at meetings or otherwise any competitively sensitive information.
 5. Don't attempt to use the Association to restrict the economic activities of any firm or any individual.
 6. Don't stay at a meeting where any such price or anti-competitive talk occurs.
 7. Do conduct all AOAC business meetings in accordance with AOAC rules. These rules require that an AOAC staff member be present or available, the meeting be conducted by a knowledgeable chair, the agenda be followed, and minutes be kept.
 8. Do confer with counsel before raising any topic or making any statement with competitive ramifications.
 9. Do send copies of meeting minutes and all AOAC-related correspondence to the staff member involved in the activity.
 10. Do alert the AOAC staff to any inaccuracies in proposed or existing methods and statements issued, or to be issued, by AOAC and to any conduct not in conformance with these guidelines.

Conclusion

Compliance with these guidelines involves not only avoidance of antitrust violations, but avoidance of any behavior which might be so construed. Bear in mind, however, that the above antitrust laws are stated in general terms, and that this statement is not a summary of applicable laws. It is intended only to highlight and emphasize the principal antitrust standards which are relevant to AOAC programs. You must, therefore, seek the guidance of either AOAC counsel or your own counsel if antitrust questions arise.

Adopted by the AOAC Board of Directors: September 24, 1989

Revised: March 11, 1991

Revised October 1996

Appendix V

POLICY ON THE USE OF THE ASSOCIATION NAME, INITIALS, IDENTIFYING INSIGNIA, LETTERHEAD, AND BUSINESS CARDS

Introduction

The following policy and guidelines for the use of the name, initials, and other identifying insignia of AOAC INTERNATIONAL have been developed in order to protect the reputation, image, legal integrity and property of the Association.

The name of the Association, as stated in its bylaws, is "AOAC INTERNATIONAL". The Association is also known by its initials, AOAC, and by its logo, illustrated below, which incorporates the Association name and a representation of a microscope, book, and flask. The AOAC logo is owned by the Association and is registered with the U.S. Patent and Trademark Office.



The full Association insignia, illustrated below, is comprised of the logo and the tagline, "The Scientific Association Dedicated to Analytical Excellence," shown below. The typeface used is Largo. The AOAC tagline is owned by the Association and is registered with the U.S. Patent and Trademark office.



The Scientific Association Dedicated to Analytical Excellence®

Policy

Policy on the use of the Association's name and logo is established by the AOAC Board of Directors as follows:

“The Board approves and encourages reference to the Association by name, either as AOAC INTERNATIONAL or as AOAC; or reference to our registered trademark, AOAC®, in appropriate settings to describe our programs, products, etc., in scientific literature and other instances so long as the reference is fair, accurate, complete and truthful and does not indicate or imply unauthorized endorsement of any kind.

The insignia (logo) of AOAC INTERNATIONAL is a registered trade and service mark and shall not be reproduced or used by any person or organization other than the Association, its elected and appointed officers, sections, or committees, without the prior written permission of the Association. Those authorized to use the AOAC INTERNATIONAL insignia shall use it only for

the purposes for which permission has been specifically granted.

The name and insignia of the Association shall not be used by any person or organization in any way which indicates, tends to indicate, or implies AOAC official endorsement of any product, service, program, company, organization, event or person, endorsement of which, has not been authorized by the Association, or which suggests that membership in the Association is available to any organization.”

The Executive Director, in accordance with the above stated policy, is authorized to process, approve, fix rules, and make available materials containing the Association name and insignia.

It should be noted that neither the Association's name nor its insignia nor part of its insignia may be incorporated into any personal, company, organization, or any other stationery other than that of the Association; nor may any statement be included in the printed portion of such stationery which states or implies that an individual, company, or other organization is a member of the Association.

Instructions

1. Reproduction or use of the Association name or insignia requires prior approval by the Executive Director or his designate.
2. Association insignia should not be altered in any manner without approval of the Executive Director or his designate, except to be enlarged or reduced in their entirety.
3. Artwork for reproducing the Association name or insignia, including those incorporating approved alterations, will be provided on request to those authorized to use them (make such requests to the AOAC Marketing Department). Examples of the types of alterations that would be approved are inclusion of a section name in or the addition of an officer's name and address to the letterhead insignia.
4. When the Association name is used without other text as a heading, it should, when possible, be set in the Largo typeface.
5. Although other colors may be used, AOAC blue, PMS 287, is the preferred color when printing the AOAC insignia, especially in formal and official documents. It is, of course, often necessary and acceptable to reproduce the insignia in black.
6. Do not print one part of the logo or insignia in one color and other parts in another color.
7. The letterhead of AOAC INTERNATIONAL shall not be used by any person or organization other than the Association, elected and appointed officers, staff, sections, or committees; except by special permission.

Correspondence of AOAC official business should be conducted using AOAC letterhead. However, those authorized to use AOAC letterhead shall use it for official AOAC business only.

Copies of all correspondence using AOAC letterhead or conducting AOAC official business,

whether on AOAC letterhead or not, must be sent to the appropriate office at AOAC headquarters.

8. AOAC INTERNATIONAL business cards shall not be used by any person or organization other than the Association, its staff, and elected officials, except by special permission.

Those authorized to use AOAC business cards shall use them for official AOAC business only and shall not represent themselves as having authority to bind the Association beyond that authorized.

Sanctions

1. Upon learning of any violation of the above policy, the Executive Director or a designate will notify the individual or organization that they are in violation of AOAC policy and will ask them to refrain from further misuse of the AOAC name or insignia.
2. If the misuse is by an Individual Member or Sustaining Member of the Association, and the misuse continues after notification, the Board of Directors will take appropriate action.
3. If continued misuse is by a nonmember of the Association or if a member continues misuse in spite of notification and Board action, ultimately, the Association will take legal action to protect its property, legal integrity, reputation, and image.

* * * * *

Adopted by the AOAC Board of Directors: September 24, 1989

Revised: June 13, 1991; February 26, 1992; March 21, 1995; October 1996



AOAC Stakeholder Panel Voting Members

AOAC INTERNATIONAL (AOAC) assembles stakeholder panels to develop voluntary consensus standards. While AOAC maintains transparency and openness in accordance with national and international guidance and regulations for standards development and its policies and procedures for assembling stakeholder panels, its policies and procedures also ensure that there is a balance of interests and perspectives in achieving consensus of the stakeholder panel.

Due Process and Balance

All AOAC stakeholder panels are diverse and can vary in size. Where a stakeholder panel is not balanced or if it is significantly large whereby consensus of the general assembly may be impractical, a balanced representative voting panel will be used to demonstrate consensus. AOAC encourages ALL stakeholders to participate in deliberations during stakeholder panel meetings and working group meetings, in addition to participating during any posted comment periods. To ensure that there is a balance of interests and perspectives, a **representative subset** of the stakeholder panel, the voting members, is selected to reach consensus for the development of AOAC voluntary consensus standards.

Composition

Voting members represent the perspectives of the larger stakeholder panel. The voting members consist of no more than $\frac{1}{4}$ to $\frac{1}{3}$ of the total number of stakeholders in registered. Primary and secondary representative voting members are approved. Every attempt is made to approve a panel of voting members that represents all perspectives of the stakeholder panel. In the event of a primary voting member is not able to attend, and no alternate has been approved, the stakeholder panel chair, working

with AOAC can provisionally approve an alternate from those in attendance to assure balance and lack of dominance. For stakeholder panels with scopes including diverse topics, the voting member representatives may be rotated to include other stakeholders for successive meetings to ensure a lack of dominance by any particular stakeholder.

Approval Process

AOAC works with the chair of the stakeholder panel and potentially other key stakeholders to develop a proposed representative voting member panel. Following AOAC policies and procedures, the proposed voting members and documentation are submitted to the AOAC Official Methods Board (OMB) for review and approval. The OMB's review ensures that the proposed panel is balanced in interests and perspectives representing the stakeholder panel and a lack of dominance.

Roles and Responsibilities

Every stakeholder has a voice and every stakeholder is entitled to state his/her or organizational perspective(s). This is due process. In developing AOAC standards, stakeholder consensus is demonstrated by 2/3 vote (67%) in favor of a motion to adopt a standard. It is important to note: Individual voting members do not have any additional weight, voice or status in stakeholder deliberations than other stakeholders. The role of the voting members is to demonstrate the consensus of the stakeholder panel. Voting members may vote in favor or against any motion and/or they may abstain. Stakeholder panel chair will moderate voting process. AOAC carefully documents the vote. It is important for voting members to be in the room during the time for voting. It is also important for voting members to inform the chair of his/her inability to serve as a voting member.



AOAC INTERNATIONAL

STAKEHOLDER PANEL ON DIETARY SUPPLEMENTS (SPDS) WORKING GROUP CHAIR & MEMBERS

VOLUNTEER ROLE DESCRIPTION

POSITION TITLE:	Working Group Chair and Members, AOAC SPDS Working Group
POSITION CLASSIFICATION:	Volunteer
REPORTS TO:	SPDS Chair
DATE PREPARED:	March 13, 2014

POSITION SUMMARY:

In keeping with the mission of AOAC INTERNATIONAL and the goals of the Stakeholder Panel on Dietary Supplements (SPDS), working group chairs will lead their working group in the development of standards (or other tasks as assigned by the SPDS chair) for specific priority ingredients as defined by the SPDS Advisory Panel. Working group chair(s) will work with AOAC staff and stakeholders to meet the working group's goals and disseminate recommendations to the stakeholder panel and community at-large. The working group may hold meetings in person and/or via teleconference (web and video) to complete its work. The chair of the working group will moderate the working group discussions, assist in scheduling the meetings, and report the working group's recommendation back to SPDS. Working group chairs will work with AOAC to formulate the working group's recommendations into motions for SPDS's consideration.

ELIGIBILITY CRITERIA FOR SPDS WORKING GROUP

CHAIR:

- Must be a key expert and/or thought leader in dietary supplements and the technologies used for priority ingredients as assigned for the specific working group.
- Must have the recommendation of the SPDS Chair.

WORKING GROUP CHAIR RESPONSIBILITIES:

- Chair meetings of the working group, moderate discussions of the working group and work with AOAC staff to facilitate working group's work.

- Work with AOAC staff and SPDS chair to identify working group members, any additional expertise/resources needed facilitate the work of the working group.
- Work as a team member and also independently.
- Present an overview on the specified priority ingredient under consideration including, but not limited to, regulatory implications, and public health and public safety challenges with methodology.
- Prepare a draft fitness for purpose statement for specified priority ingredient and technology to present to SPDS for consideration.
- Work with AOAC staff to reconcile actions and outcomes of working group deliberations.
- Using AOAC guidance to reconcile comments and address questions on SMPR.
- Present working group recommended SMPR to SPDS for review and approval.
- Work with AOAC staff and stakeholders to draft and review relevant methodology and working group documentation.
- Draft SMPR white paper for publication.
- Perform duties and reviews in timely fashion.
- Other tasks as agreed upon by working group chair, SPDS chair and AOAC staff.

DUTIES AND RESPONSIBILITIES OF THE SPDS WORKING GROUP MEMBERS:

The working group will meet either in person and via teleconference, web conferencing or by other means of communication. All communication and meetings of the working group must be facilitated through AOAC

staff. The working group's tasks will include developing standard method performance requirements (SMPRs), review of methodology, identifying expertise and other as may be requested by the SPDS chair. Working groups are not required to vote, but to show general consensus for its recommendations. The groups should meet to discuss their objectives and complete their assigned tasks. Individuals on the working groups may be tasked with their own action items and responsibilities. More than one meeting and one round of communication may be required to complete the working group's tasks. All working group participants are expected to contribute and are expected to have completed the SMPR Education Session. AOAC staff will document all working group decisions and actions.

AOAC RESOURCES:

- Referencing AOAC guidance documentation to assist in drafting the fitness for purpose statement,

standard method performance requirements (SMPR), and additional work as tasked.

- 1) AOAC Fitness for Purpose Statement Guideline
- 2) Appendix F: Guidelines for Standard Method Performance Requirements
- 3) Appendix K: Guidelines for Dietary Supplements and Botanicals

STAFF LIASON:

AOAC will assign staff to facilitate the work of the working group.

TERMS OF REVIEW:

This document will be reviewed biannually by the SPDS Chair and AOAC staff.

DATES REVISED:

Stakeholder Panel

Voting Panel – A vetted, representative, and balanced subset of the assembled stakeholders. Ideally the number of voters represents $\frac{1}{4}$ to $\frac{1}{3}$ of the assembly.

Voting Guidelines – A. motions to create a consensus based standard (ex: voting on fitness for purpose statements or Standard Method Performance Requirements) require a 2/3 vote for the motion to carry.
 B. Any other motion (ex: votes to clarify information for working groups, set priorities or direction, etc.) requires a majority vote to carry.

Expert Review Panel

Voting Panel – 7 – 10 vetted experts

Quorum - The presence of **7** members or **2/3** of total vetted ERP membership, whichever is greater.

Voting Guidelines – Motions to adopt a *First Action Official MethodSM of Analysis* carry by unanimous vote on first ballot. If not unanimous, negative votes must delineate scientific reasons, and can be overridden by 2/3 of voting ERP members after due consideration. Dissenting opinions are recorded.

Working Group

Voting Panel – There is no formal voting panel. Any interested and knowledgeable party may participate. Working groups sole purpose is to provide recommendations to stakeholder panels.

Voting Guidelines –majority vote carries all motions, dissenting opinions considered by assembly and recorded.

Helpful Definitions & Terminology

Quorum	The number of members who must be present in order to validly transact business. It is determined by the number of members present, not the number present and voting. <i>(Fundamentals of Parliamentary Law and Procedure, 3rd edition. p. 151).</i>
Representative Voting Panel Members	Every member has an obligation to vote and the right to abstain.
Abstentions	Abstentions reduce the number required to obtain a majority of those present and voting. They are only counted to confirm the presence of a quorum. <i>(Fundamentals of Parliamentary Law and Procedure, 3rd edition. p. 237).</i>
Order	Meetings should address only one item of business at one time (only one pending motion at a time). Chairs should not permit digression or introduction of different topics until the business at hand is resolved. No pending motions while changing topics. <i>(Fundamentals of Parliamentary Law and Procedure, 3rd edition. p. 1).</i> All business must be conducted with order and should be done fairly and impartially. The presiding officer should impartially ensure that each member has an opportunity to speak. <i>(Fundamentals of Parliamentary Law and Procedure, 3rd edition. pp. 1-2).</i>
Equality	All members have equal opportunity to propose motions, to participate in debate, to vote, to serve on committees or as an officer, to share in activities according to the member’s abilities. <i>(Fundamentals of Parliamentary Law and Procedure, 3rd edition. p. 2).</i>
Justice	All members have the right to ask questions, to be informed, to have complex motions explained by the chair. <i>(Fundamentals of Parliamentary Law and Procedure, 3rd edition. p. 2).</i>
Minority Rights	Dissenting members have equal rights to voice opposing or minority opinions and strive to become the majority. <i>(Fundamentals of Parliamentary Law and Procedure, 3rd edition. p. 2).</i>
Majority Rights	No members, board, or officers have the right to dictate or control decisions unless the member grant such rights Members may not take any action in conflict with federal, regional or organizational laws or policies. Decisions are based on the will of the majority. <i>(Fundamentals of Parliamentary Law and Procedure, 3rd edition. p. 2).</i>

Appendix F: Guidelines for Standard Method Performance Requirements

Contents

Introduction to Standard Method Performance Requirements	1
Annex A: Format of a Standard Method Performance Requirement	5
Annex B: Classification of Methods	11
Annex C: Understanding the POD Model	12
Annex D: Definitions and Calculations of HorRat Values from Intralaboratory Data	13
Annex E: AOAC Method Accuracy Review	15
Annex F: Development and Use of In-House Reference Materials	16

Introduction to Standard Method Performance Requirements

Standard method performance requirements (SMPRs) are a unique and novel concept for the analytical methods community. SMPRs are voluntary consensus standards, developed by stakeholders, that prescribe the minimum analytical performance requirements for classes of analytical methods. In the past, analytical methods were evaluated and the results compared to a “gold standard” method, or if a gold standard method did not exist, then reviewers would decide retrospectively if the analytical performance was acceptable. Frequently, method developers concentrated on the process of evaluating the performance parameters of a method, and rarely set acceptance criteria. However, as the *Eurachem Guide* points out: “. . . the judgment of method suitability for its intended use is equally important . . .” (1) to the evaluation process.

International Voluntary Consensus Standards

An SMPR is a form of an international, voluntary consensus standard. A standard is an agreed, repeatable way of doing something that is published as document that contains a technical specification or other precise criteria designed to be used consistently as a rule, guideline, or definition. SMPRs are a *consensus* standards developed by stakeholders in a very controlled process that ensures that users, research organizations, government departments, and consumers work together to create a standard that meets the demands of the analytical community and technology. SMPRs are also *voluntary* standards. AOAC cannot, and does not, impose the use of SMPRs. Users are free to use SMPRs as they see fit. AOAC is very careful to include participants from as many regions of the world as possible so that SMPRs are accepted as *international* standards.

Guidance for Standard Method Performance Requirements

Commonly known as the “SMPR Guidelines.” The first version of the SMPR Guidelines were drafted in 2010 in response to the increasing use and popularity of SMPRs as a vehicle to describe the analytical requirements of a method. Several early “acceptance

criteria” documents were prepared for publication in late 2009, but the format of the acceptance criteria documents diverged significantly from one another in basic format. AOAC realized that a guidance document was needed to promote uniformity.

An early version of the SMPR Guidelines were used for a project to define the analytical requirements for endocrine disruptors in potable water. The guidelines proved to be extremely useful in guiding the work of the experts and resulted in uniform SMPRs. Subsequent versions of the SMPR Guidelines were used in the Stakeholder Panel for Infant Formula and Adult Nutritionals (SPIFAN) project with very positive results. The SMPR Guidelines are now published for the first time in the *Journal of AOAC INTERNATIONAL* and *Official Methods of Analysis*.

Users of the guidelines are advised that they are: (1) a *guidance* document, not a statute that users must conform to; and (2) a “living” document that is regularly updated, so users should check the AOAC website for the latest version before using these guidelines.

The SMPR Guidelines are intended to provide basic information for working groups assigned to prepare SMPRs. The guidelines consist of the standard format of an SMPR, followed by a series of informative tables and annexes.

SMPR Format

The general format for an SMPR is provided in *Annex A*.

Each SMPR is identified by a unique SMPR number consisting of the year followed by a sequential identification number (YYYY.XXX). An SMPR number is assigned when the standard is approved. By convention, the SMPR number indicates the year a standard is approved (as opposed to the year the standard is initiated). For example, SMPR 2010.003 indicates the third SMPR adopted in 2010.

The SMPR number is followed by a method name that must include the analyte(s), matrix(es), and analytical technique (unless the SMPR is truly intended to be independent of the analytical technology). The method name may also refer to a “common” name (e.g., “Kjeldahl” method).

The SMPR number and method name are followed by the name of the stakeholder panel or expert review panel that approved the SMPR, and the approval and effective dates.

Information about method requirements is itemized into nine categories: (1) intended use; (2) applicability; (3) analytical technique; (4) definitions; (5) method performance requirements; (6) system suitability; (7) reference materials; (8) validation guidance; and (9) maximum time-to-determination.

An SMPR for qualitative and/or identification methods may include up to three additional annexes: (1) inclusivity/selectivity panel; (2) exclusivity/cross-reactivity panel; and (3) environmental material panels. These annexes not required.

Informative tables.—The SMPR Guidelines contain seven informative tables that represent the distilled knowledge of many years of method evaluation, and are intended as guidance for SMPR working groups. The informative tables are not necessarily AOAC

policy. SMPR working groups are expected to apply their expertise in the development of SMPRs.

Table A1: Performance Requirements. Provides recommended performance parameters to be included into an SMPR. Table A1 is organized by five method classifications: (1) main component quantitative methods; (2) trace or contaminant quantitative methods; (3) main component qualitative methods; (4) trace or contaminant quantitative methods; and (5) identification methods. The table is designed to accommodate both microbiological and chemical methods. Alternate microbiological/chemical terms are provided for equivalent concepts.

Table A2: Recommended Definitions. Provides definitions for standard terms in the SMPR Guidelines. AOAC relies on *The International Vocabulary of Metrology Basic and General Concepts and Associated Terms* (VIM) and the International Organization for Standardization (ISO) for definition of terms not included in Table A2.

Table A3: Recommendations for Evaluation. Provides general guidance for evaluation of performance parameters. More detailed evaluation guidance can be found in *Appendix D, Guidelines for Collaborative Study Procedures to Validate Characteristics of a Method of Analysis* (2); *Appendix I, Guidelines for Validation of Biological Threat Agent Methods and/or Procedures* (3); *Appendix K, AOAC Guidelines for Single-Laboratory Validation of Chemical Methods for Dietary Supplements and Botanicals* (4); *Codex Alimentarius Codex Procedure Manual* (5); and *ISO Standard 5725-1-1994* (6).

Table A4: Expected Precision (Repeatability) as a Function of Analyte Concentration. The precision of a method is the closeness of agreement between independent test results obtained under stipulated conditions. Precision is usually expressed in terms

of imprecision and computed as a relative standard deviation (RSD) of the test results. The imprecision of a method increases as the concentration of the analyte decreases. This table provides target RSDs for a range of analyte concentrations.

Table A5: Expected Recovery as a Function of Analyte Concentration. Recovery is defined as the ratio of the observed mean test result to the true value. The range of the acceptable mean recovery expands as the concentration of the analyte decreases. This table provides target mean recovery ranges for analyte concentrations from 1 ppb to 100%.

Table A6: Predicted Relative Standard Deviation of Reproducibility (PRSD_R). This table provides the calculated PRSD_R using the Horwitz formula:

$$PRSD_R = 2C^{-0.15}$$

where C is expressed as a mass fraction.

Table A7: POD and Number of Test Portions. This table provides the calculated probability of detection (POD) for given sample sizes and events (detections). A method developer can use this table to determine the number of analyses required to obtain a specific POD.

Informative annexes.—The SMPR Guidelines contain informative annexes on the topics of classification of methods, POD model, HorRat values, reference materials, and method accuracy and review. As with the informative tables, these annexes are intended to provide guidance and information to the working groups.

Initiation of an SMPR

See Figure 1 for a schematic flowchart diagram of the SMPR development process.




Figure 1. Schematic flowchart diagram of the SMPR development process.

Advisory panels.—Most commonly, an SMPR is created in response to an analytical need identified by an advisory panel. Advisory panels normally consist of sponsors and key stakeholders who have organized to address analytical problems. Usually, the advisory panel identifies general analytical problems, such as the need to update analytical methods for determination of nutrients in infant formula. An advisory panel, with the input of appropriate subject matter experts, also prioritizes the specific analytical problems within the general topic. This panel is critical in planning for the stakeholder panel meeting.

Stakeholder panels.—After an advisory panel has identified a general analytical problem, AOAC announces the standards development activity, identifies stakeholders, and organizes a stakeholder panel. Membership on a stakeholder panel is open to anyone materially affected by the proposed standard. AOAC recruits scientists to participate on stakeholder panels on the basis of their expertise with the analytical problem identified by the advisory panel. Experts are recruited from academia, government, nongovernmental organizations (such as ISO), industry, contract research organizations, method developers, and instrument/equipment manufacturers. AOAC employs a representative voting panel model to ensure balance with regards to stakeholder perspective, and to ensure that no particular stakeholder perspective dominates the proceedings of the stakeholder panel. All stakeholder candidates are reviewed by the AOAC Chief Scientific Officer (CSO) for relevant qualifications, and again by the Official Methods Board to ensure that the stakeholder panel is balanced and all stakeholders are fairly represented.

Stakeholder panels are extremely important as they serve several functions: (1) identify specific analytical topics within the general analytical problem described by the advisory panel; (2) form working groups to address the specific analytical topics; (3) identify additional subject matter experts needed for the working groups; (4) provide oversight of the SMPR development; and (5) formally adopt SMPRs originally drafted by working groups.

Working groups.—Working groups are formed by the stakeholder panel when a specific analytical topic has been identified. The primary purpose of a working group is to draft an SMPR. Working groups may also be formed to make general recommendations, such as developing a common definition to be used by multiple working groups. For example, SPIFAN formed a working group to create a definition for “infant formula” that could be shared and used by all of the SPIFAN working groups.

The process of drafting an SMPR usually requires several months, and several meetings and conference calls. An SMPR drafted by a working group is presented to a stakeholder panel. A stakeholder panel may revise, amend, or adopt a proposed SMPR on behalf of AOAC.

Fitness-for-Purpose Statement and Call for Methods

One of the first steps in organizing a project is creating a fitness-for-purpose statement. In AOAC, the fitness-for-purpose statement is a very general description of the methods needed. It is the responsibility of a working group chair to draft a fitness-for-purpose statement. A working group chair is also asked to prepare a presentation with background information about the analyte, matrix, and the nature of the analytical problem. A working group chair presents the background information and proposes a draft fitness-for-purpose statement to the presiding stakeholder panel. The stakeholder panel is asked to endorse the fitness-for-purpose statement.

The AOAC CSO prepares a call for methods based on the stakeholder panel-approved fitness-for-purpose statement. The call for methods is posted on the AOAC website and/or e-mailed to the AOAC membership and other known interested parties. AOAC staff collects and compiles candidate methods submitted in response to the call for methods. The CSO reviews and categorizes the methods.

Creating an SMPR

Starting the process of developing an SMPR can be a daunting challenge. In fact, drafting an SMPR should be a daunting challenge because the advisory panel has specifically identified an analytical problem that has yet to be resolved. Completing an SMPR can be a very rewarding experience because working group members will have worked with their colleagues through a tangle of problems and reached a consensus where before there were only questions.

It is advisable to have some representative candidate methods available for reference when a working group starts to develop an SMPR. These methods may have been submitted in response to the call for methods, or may be known to a working group member. In any case, whatever the origin of the method, candidate methods may assist working group members to determine reasonable performance requirements to be specified in the SMPR. The performance capabilities of existing analytical methodologies is a common question facing a working group.

Normally, a working chair and/or the AOAC CSO prepares a draft SMPR. A draft SMPR greatly facilitates the process and provides the working group with a structure from which to work.

Working group members are advised to first consider the “intended use” and “maximum time-to-determination” sections as this will greatly affect expectations for candidate methods. For example, methods intended to be used for surveillance probably need to be quick but do not require a great deal of precision, and false-positive results might be more tolerable. Whereas methods intended to be used for dispute resolution will require better accuracy, precision, and reproducibility, but time to determination is not as important.

Once a working group has agreed on the intended use of candidate methods, then it can begin to define the applicability of candidate methods. The applicability section of the SMPR is one of the most important, and sometimes most difficult, sections of the SMPR. The analyte(s) and matrix(es) must be explicitly identified. For chemical analytes, International Union of Pure and Applied Chemistry (IUPAC) nomenclature and/or Chemical Abstracts Service (CAS) registry numbers should be specified. Matrix(es) should be clearly identified including the form of the matrix such as raw, cooked, tablets, powders, etc. The nature of the matrix may affect the specific analyte. It may be advantageous to fully identify and describe the matrix before determining the specific analyte(s). It is not uncommon for working groups to revise the initial definition of the analyte(s) after the matrix(es) has been better defined.

Table 1. Example of method performance table for a single analyte

Analytical range	7.0–382.6 µg/mL	
Limit of quantitation (LOQ)	≤7.0 µg/mL	
Repeatability (RSD,)	<10 µg/mL	≤8%
	≥10 µg/mL	≤6%

Table 2. Example of method performance table for multiple analytes

	Analyte 1		Analyte 2		Analyte 3	
Analytical range	10–20 µg/mL		100–200 µg/mL		200–500 µg/mL	
Limit of quantitation (LOQ)	≤10 µg/mL		≤100 µg/mL		≤200 µg/mL	
Repeatability (RSD,)	<10 µg/mL	≤8%	<10 µg/mL	≤8%	<200 µg/mL	≤10%
	≥10 µg/mL	≤6%	≥10 µg/mL	≤6%	≥200 µg/mL	≤8%

For projects with multiple analytes, for example, vitamins A, D, E, and K in infant formula, it may be useful to organize a separate working group to fully describe the matrix(es) so that a common description of the matrix(es) can be applied to all of the analytes.

For single analyte SMPRs, it is most common to organize the method performance requirements into a table with 2–3 columns as illustrated in Table 1. For multiple analyte SMPRs, it is often convenient to present the requirements in an expanded table with analytes forming additional columns as illustrated in Table 2.

Once the intended use, analytical techniques, and method performance requirements have been determined, then a working group can proceed to consider the quality control parameters, such as the minimum validation requirements, system suitability procedures, and reference materials (if available). It is not uncommon that an appropriate reference material is not available. *Annex F* of the SMPR Guidelines provides comprehensive guidance for the development and use of in-house reference materials.

Most working groups are able to prepare a consensus SMPR in about 3 months.

Open Comment Period

Once a working group has produced a draft standard, AOAC opens a comment period for the standard. The comment period provides an opportunity for other stakeholders to state their perspective on the draft SMPR. All collected comments are reviewed by the AOAC CSO and the working group chair, and the comments are reconciled. If there are significant changes required to the draft standard as a result of the comments, the working group is convened to discuss and any unresolved issues will be presented for discussion at the stakeholder panel meeting.

Submission of Draft SMPRs to the Stakeholder Panel

Stakeholder panels meet several times a year at various locations. The working group chair (or designee) presents a draft SMPR to the stakeholder panel for review and discussion. A working group chair is expected to be able to explain the conclusions of the working group, discuss comments received, and to answer questions from the stakeholder panel. The members of the stakeholder panel may revise, amend, approve, or defer a decision on the proposed SMPR. A super majority of 2/3 or more of those voting is required to adopt an SMPR as an AOAC voluntary consensus standard.

Publication

Adopted SMPRs are prepared for publication by AOAC staff, and are published in the *Journal of AOAC INTERNATIONAL* and in the AOAC *Official Methods of Analysis*SM compendium. Often, the AOAC CSO and working group chair prepare a companion article to introduce an SMPR and describe the analytical issues considered and resolved by the SMPR. An SMPR is usually published within 6 months of adoption.

Conclusion

SMPRs are a unique and novel concept for the analytical methods community. SMPRs are voluntary, consensus standards developed by stakeholders that prescribe the minimum analytical performance requirements for classes of analytical methods. The SMPR Guidelines provide a structure for working groups to use as they develop an SMPR. The guidelines have been employed in several AOAC projects and have been proven to be very useful. The guidelines are not a statute that users must conform to; they are a “living” document that is regularly updated, so users should check the AOAC website for the latest version before using the guidelines.

References

- (1) Eurachem, *The Fitness for Purpose of Analytical Methods: A Laboratory Guide to Method Validation and Related Topics, Validation*, <http://www.eurachem.org/guides/pdf/valid.pdf>, posted December 1998, accessed March 2012
- (2) *Guidelines for Collaborative Study Procedures to Validate Characteristics of a Method of Analysis* (2012) *Official Methods of Analysis, Appendix D*, AOAC INTERNATIONAL, Gaithersburg, MD
- (3) *AOAC INTERNATIONAL Methods Committee Guidelines for Validation of Biological Threat Agent Methods and/or Procedures* (2012) *Official Methods of Analysis*, 19th Ed., *Appendix I, Calculation of CPOD and dCPOD Values from Qualitative Method Collaborative Study Data*, AOAC INTERNATIONAL, Gaithersburg, MD
- (4) *AOAC Guidelines for Single-Laboratory Validation of Chemical Methods for Dietary Supplements and Botanicals* (2012) *Official Methods of Analysis*, 19th Ed., *Appendix K*, AOAC INTERNATIONAL, Gaithersburg, MD
- (5) Codex Alimentarius Codex Procedure Manual
- (6) International Organization for Standardization, Geneva, Switzerland

ANNEX A
Format of a
Standard Method Performance Requirement

AOAC SMPR YYYY.XXX
(YYYY = Year; XXX = sequential identification number)

Method Name: Must include the analyte(s), matrix(es), and analytical technique [unless the standard method performance requirement (SMPR) is truly intended to be independent of the analytical technology]. The method name may refer to a “common” name (e.g., “Kjeldahl” method).

Approved By: Name of stakeholder panel or expert review panel

Final Version Date: Date

Effective Date: Date

1. Intended Use: Additional information about the method and conditions for use.

2. Applicability: List matrixes if more than one. Provide details on matrix such as specific species for biological analytes, or International Union of Pure and Applied Chemistry (IUPAC) nomenclature and Chemical Abstracts Service (CAS) registry number for chemical analytes. Specify the form of the matrix such as raw, cooked, tablets, powders, etc.

3. Analytical Technique: Provide a detailed description of the analytical technique if the SMPR is to apply to a specific analytical technique; or state that the SMPR applies to any method that meets the method performance requirements.

4. Definitions: List and define terms used in the performance parameter table (*see* Table A2 for list of standard terms).

5. Method Performance Requirements: List the performance parameters and acceptance criteria appropriate for each method/analyte/matrix. *See* Table A1 for appropriate performance requirements.

If more than one analyte/matrix, and if acceptance criteria differ for analyte/matrix combinations then organize a table listing each analyte/matrix combination and its minimum acceptance criteria for each performance criteria.

6. System Suitability Tests and/or Analytical Quality Control: Describe minimum system controls and QC procedures.

7. Reference Material(s): Identify the appropriate reference materials if they exist, or state that reference materials are not available. Refer to *Annex E (AOAC Method Accuracy Review)* for instructions on the use of reference materials in evaluations.

8. Validation Guidance: Recommendations for type of evaluation or validation program such as single-laboratory validation (SLV), *Official Methods of Analysis*SM (OMA), or *Performance Tested Methods*SM (PTM).

9. Maximum Time-to-Determination: Maximum allowable time to complete an analysis starting from the test portion preparation to final determination or measurement.

Annex I: Inclusivity/Selectivity Panel. Recommended for qualitative and identification method SMPRs.

Annex II: Exclusivity/Cross-Reactivity Panel. Recommended for qualitative and identification method SMPRs.

Annex III: Environmental Materials Panel. Recommended for qualitative and identification method SMPRs.

Table A1. Performance requirements

Classifications of methods ^a				
Quantitative method		Qualitative method		Identification method
Main component ^b	Trace or contaminant ^c	Main component ^b	Trace or contaminant ^c	
Parameter				
Single-laboratory validation				
Applicable range	Applicable range	Inclusivity/selectivity	Inclusivity/selectivity	Inclusivity/selectivity
Bias ^d	Bias ^d	Exclusivity/cross-reactivity	Exclusivity/cross-reactivity	Exclusivity/cross-reactivity
Precision	Precision	Environmental interference	Environmental interference	Environmental interference
Recovery	Recovery	Laboratory variance	Laboratory variance	
Limit of quantitation (LOQ)	LOQ	Probability of detection (POD) ^e	POD at AMDL ^f	Probability of identification (POI)
Reproducibility				
RSD _r or target measurement uncertainty	RSD _r or target measurement uncertainty	POD (0) POD (c) Laboratory POD ^g	POD (0) POD (c) Laboratory POD ^g	POI (c) Laboratory POI

^a See Annex B for additional information on classification of methods.

^b ≥100 g/kg.

^c <100 g/kg.

^d If a reference material is available.

^e At a critical level.

^f AMDL = Acceptable minimum detection level.

^g LPOD = CPOD.

Table A2. Recommended definitions

Bias	Difference between the expectation of the test results and an accepted reference value. Bias is the total systematic error as contrasted to random error. There may be one or more systematic error components contributing to the bias.
Environmental interference	Ability of the assay to detect target organism in the presence of environmental substances and to be free of cross reaction from environmental substances.
Exclusivity	Strains or isolates or variants of the target agent(s) that the method must not detect.
Inclusivity	Strains or isolates or variants of the target agent(s) that the method can detect.
Laboratory probability of detection (POD)	Overall fractional response (mean POD = CPOD) for the method calculated from the pooled POD_j responses of the individual laboratories ($j = 1, 2, \dots, L$). ^a See Annex C.
Limit of quantitation (LOQ)	Minimum concentration or mass of analyte in a given matrix that can be reported as a quantitative result.
POD (0)	Probability of the method giving a (+) response when the sample is truly without analyte.
POD (c)	Probability of the method giving a (–) response when the sample is truly without analyte.
POD	Proportion of positive analytical outcomes for a qualitative method for a given matrix at a given analyte level or concentration. Consult Annex C for a full explanation.
Probability of identification (POI)	Expected or observed fraction of test portions at a given concentration that gives positive result when tested at a given concentration. Consult <i>Probability of Identification (POI): A Statistical Model for the Validation of Qualitative Botanical Identification Methods</i> . ^c
Precision (repeatability)	Closeness of agreement between independent test results obtained under stipulated conditions. The measure of precision is usually expressed in terms of imprecision and computed as a standard deviation of the test results. ^d
Recovery	Fraction or percentage of the analyte that is recovered when the test sample is analyzed using the entire method. There are two types of recovery: (1) Total recovery based on recovery of the native plus added analyte, and (2) marginal recovery based only on the added analyte (the native analyte is subtracted from both the numerator and denominator). ^e
Repeatability	Precision under repeatability conditions.
Repeatability conditions	Conditions where independent test results are obtained with the same method on identical test items in the same laboratory by the same operator using the same equipment within short intervals of time.
Reproducibility	Precision under reproducibility conditions.
Reproducibility conditions	Conditions where independent test results are obtained with the same method on identical test items in different laboratories with different operators using different equipment.
Relative standard deviation (RSD)	$RSD = s_i \times 100/\bar{x}$
Standard deviation (s_i)	$s_i = [\sum(x_i - \bar{x})^2/n]^{0.5}$

^a AOAC INTERNATIONAL Methods Committee Guidelines for Validation of Biological Threat Agent Methods and/or Procedures (Calculation of CPOD and dCPOD Values from Qualitative Method Collaborative Study Data), *J. AOAC Int.* **94**, 1359(2011) and *Official Methods of Analysis of AOAC INTERNATIONAL* (2012) 19th Ed., Appendix I.

^b *International Vocabulary of Metrology (VIM)—Basic and General Concepts and Associated Terms* (2008) JCGM 200:2008, Joint Committee for Guides in Metrology (JCGM), www.bipm.org

^c LaBudde, R.A., & Harnly, J.M. (2012) *J. AOAC Int.* **95**, 273–285.

^d ISO 5725-1-1994.

^e *Official Methods of Analysis* (2012) Appendix D (Guidelines for Collaborative Study Procedures to Validate Characteristics of a Method of Analysis), AOAC INTERNATIONAL, Gaithersburg, MD.

Table A3. Recommendations for evaluation

Bias (if a reference material is available)	A minimum of five replicate analyses of a Certified Reference Material. ^a
Environmental interference	Analyze test portions containing a specified concentration of one environmental materials panel member. Materials may be pooled. Consult with AOAC statistician.
Exclusivity/cross-reactivity	Analyze one test portion containing a specified concentration of one exclusivity panel member. More replicates can be used. Consult with AOAC statistician.
Inclusivity/selectivity	Analyze one test portion containing a specified concentration of one inclusivity panel member. More replicates can be used. Consult with AOAC statistician.
Limit of quantitation (LOQ)	Estimate the LOQ = average (blank) + 10 × s ₀ (blank). Measure blank samples with analyte at the estimated LOQ. Calculate the mean average and standard deviation of the results. Guidance ^b : For ML ≥ 100 ppm (0.1 mg/kg): LOD = ML × 1/5. For ML < 100 ppm (0.1 mg/kg): LOD = ML × 2/5.
Measurement uncertainty	Use ISO 21748: <i>Guidance for the use of repeatability, reproducibility, and trueness estimates in measurement uncertainty estimation to analyze data collected for bias, repeatability, and intermediate precision to estimate measurement uncertainty.</i>
POD(0)	Use data from collaborative study.
POD (c)	
Repeatability	Prepare and homogenize three unknown samples at different concentrations to represent the full, claimed range of the method. Analyze each unknown sample by the candidate method seven times, beginning each analysis from weighing out the test portion through to final result with no additional replication (unless stated to do so in the method). All of the analyses for one unknown sample should be performed within as short a period of time as is allowed by the method. The second and third unknowns may be analyzed in another short time period. Repeat for each claimed matrix.
Probability of detection (POD)	Determine the desired POD at a critical concentration. Consult with Table A7 to determine the number of test portions required to demonstrate the desired POD.
Probability of identification (POI)	Consult <i>Probability of Identification (POI): A Statistical Model for the Validation of Qualitative Botanical Identification Methods</i> ^c .
Recovery	Determined from spiked blanks or samples with at least seven independent analyses per concentration level at a minimum of three concentration levels covering the analytical range. Independent means at least at different times. If no confirmed (natural) blank is available, the average inherent (naturally containing) level of the analyte should be determined on at least seven independent replicates. Marginal % recovery = $(C_f - C_u) \times 100 / C_A$ Total % recovery = $100(C_f) / (C_u + C_A)$ where C _f = concentration of fortified samples, C _u = concentration of unfortified samples, and C _A = concentration of analyte added to the test sample. ^d Usually total recovery is used unless the native analyte is present in amounts greater than about 10% of the amount added, in which case use the method of addition. ^e
Reproducibility (collaborative or interlaboratory study)	Quantitative methods: Recruit 10–12 collaborators; must have eight valid data sets; two blind duplicate replicates at five concentrations for each analyte/matrix combination to each collaborator.
	Qualitative methods: Recruit 12–15 collaborators; must have 10 valid data sets; six replicates at five concentrations for each analyte/matrix combination to each collaborator.

^a *Guidance for Industry for Bioanalytical Method Validation* (May 2001) U.S. Department of Health and Human Services, U.S. Food and Drug Administration, Center for Drug Evaluation and Research (CDER), Center for Veterinary Medicine (CVM).

^b Codex Alimentarius Codex Procedure Manual.

^c LaBudde, R.A., & Harnly, J.M. (2012) *J. AOAC Int.* **95**, 273–285.

^d *Guidelines for Collaborative Study Procedures to Validate Characteristics of a Method of Analysis* (2012) *Official Methods of Analysis*, 19th Ed., Appendix D, AOAC INTERNATIONAL, Gaithersburg, MD.

^e *AOAC Guidelines for Single-Laboratory Validation of Chemical Methods for Dietary Supplements and Botanicals* (2012) *Official Methods of Analysis*, 19th Ed., Appendix K, AOAC INTERNATIONAL, Gaithersburg, MD.

Table A4. Expected precision (repeatability) as a function of analyte concentration^a

Analyte, %	Analyte ratio	Unit	RSD, %
100	1	100%	1.3
10	10 ⁻¹	10%	1.9
1	10 ⁻²	1%	2.7
0.01	10 ⁻³	0.1%	3.7
0.001	10 ⁻⁴	100 ppm (mg/kg)	5.3
0.0001	10 ⁻⁵	10 ppm (mg/kg)	7.3
0.00001	10 ⁻⁶	1 ppm (mg/kg)	11
0.000001	10 ⁻⁷	100 ppb (µg/kg)	15
0.0000001	10 ⁻⁸	10 ppb (µg/kg)	21
0.00000001	10 ⁻⁹	1 ppb (µg/kg)	30

^a Table excerpted from AOAC Peer-Verified Methods Program, Manual on Policies and Procedures (1998) AOAC INTERNATIONAL, Gaithersburg, MD.

The precision of a method is the closeness of agreement between independent test results obtained under stipulated conditions. Precision is usually expressed in terms of imprecision and computed as a relative standard deviation of the test results. The imprecision of a method increases as the concentration of the analyte decreases. This table provides targets RSDs for a range of analyte concentrations.

Table A5. Expected recovery as a function of analyte concentration^a

Analyte, %	Analyte ratio	Unit	Mean recovery, %
100	1	100%	98–102
10	10 ⁻¹	10%	98–102
1	10 ⁻²	1%	97–103
0.01	10 ⁻³	0.1%	95–105
0.001	10 ⁻⁴	100 ppm	90–107
0.0001	10 ⁻⁵	10 ppm	80–110
0.00001	10 ⁻⁶	1 ppm	80–110
0.000001	10 ⁻⁷	100 ppb	80–110
0.0000001	10 ⁻⁸	10 ppb	60–115
0.00000001	10 ⁻⁹	1 ppb	40–120

^a Table excerpted from AOAC Peer-Verified Methods Program, Manual on Policies and Procedures (1998) AOAC INTERNATIONAL, Gaithersburg, MD.

Recovery is defined as the ratio of the observed mean test result to the true value. The range of the acceptable mean recovery expands as the concentration of the analyte decreases. This table provides target mean recovery ranges for analyte concentrations from 100% to 1 ppb.

Table A6. Predicted relative standard deviation of reproducibility (PRSD_R)^a

Concentration (C)	Mass fraction (C)	PRSD _R , %
100%	1.0	2
1%	0.01	4
0.01%	0.0001	8
1 ppm	0.000001	16
10 ppb	0.00000001	32
1 ppb	0.000000001	45

^a Table excerpted from *Definitions and Calculations of HorRat Values from Intralaboratory Data*, HorRat for SLV.doc, 2004-01-18, AOAC INTERNATIONAL, Gaithersburg, MD.

Predicted relative standard deviation = PRSD_R. Reproducibility relative standard deviation calculated from the Horwitz formula:

$$\text{PRSD}_R = 2C^{-0.15}, \text{ where } C \text{ is expressed as a mass fraction}$$

This table provides the calculated PRSD_R for a range of concentrations. See Annex D for additional information.

Table A7. POD and number of test portions^{a,b}

Sample size required for proportion							
Assume	1. Binary outcome (occur/not occur). 2. Constant probability rho of event occurring. 3. Independent trials (e.g., simple random sample). 4. Fixed number of trials (N)						
Inference	95% Confidence interval lies entirely at or above specified minimum rho						
Desired	Sample size N needed						
Minimum probability rho, %	Sample size (N)	Minimum No. events (x)	Maximum No. nonevents (y)	1-Sided lower confidence limit on rho ^c , %	Expected lower confidence limit on rho, %	Expected upper confidence limit on rho, %	Effective AOQL ^d rho, %
50	3	3	0	52.6	43.8	100.0	71.9
50	10	8	2	54.1	49.0	94.3	71.7
50	20	14	6	51.6	48.1	85.5	66.8
50	40	26	14	52.0	49.5	77.9	63.7
50	80	48	32	50.8	49.0	70.0	59.5
55	4	4	0	59.7	51.0	100.0	75.5
55	10	9	1	65.2	59.6	100.0	79.8
55	20	15	5	56.8	53.1	88.8	71.0
55	40	28	12	57.1	54.6	81.9	68.2
55	80	52	28	55.9	54.1	74.5	64.3
60	5	5	0	64.9	56.5	100.0	78.3
60	10	9	1	65.2	59.6	100.0	79.8
60	20	16	4	62.2	58.4	91.9	75.2
60	40	30	10	62.4	59.8	85.8	72.8
60	80	56	24	61.0	59.2	78.9	69.1
65	6	6	0	68.9	61.0	100.0	80.5
65	10	9	1	65.2	59.6	100.0	79.8
65	20	17	3	67.8	64.0	94.8	79.4
65	40	31	9	65.1	62.5	87.7	75.1
65	80	59	21	65.0	63.2	82.1	72.7
70	7	7	0	72.1	64.6	100.0	82.3
70	10	10	0	78.7	72.2	100.0	86.1
70	20	18	2	73.8	69.9	97.2	83.6
70	40	33	7	70.7	68.0	91.3	79.7
70	80	63	17	70.4	68.6	86.3	77.4
75	9	9	0	76.9	70.1	100.0	85.0
75	10	10	0	78.7	72.2	100.0	86.1
75	20	19	1	80.4	76.4	100.0	88.2
75	40	35	5	76.5	73.9	94.5	84.2
75	80	67	13	75.9	74.2	90.3	82.2
80	11	11	0	80.3	74.1	100.0	87.1
80	20	19	1	80.4	76.4	100.0	88.2
80	40	37	3	82.7	80.1	97.4	88.8
80	80	70	10	80.2	78.5	93.1	85.8
85	20	20	0	88.1	83.9	100.0	91.9
85	40	38	2	86.0	83.5	98.6	91.1
85	80	74	6	86.1	84.6	96.5	90.6
90	40	40	0	93.7	91.2	100.0	95.6
90	60	58	2	90.4	88.6	99.1	93.9
90	80	77	3	91.0	89.5	98.7	94.1
95	60	60	0	95.7	94.0	100.0	97.0
95	80	80	0	96.7	95.4	100.0	97.7
95	90	89	1	95.2	94.0	100.0	97.0
95	96	95	1	95.5	94.3	100.0	97.2
98	130	130	0	98.0	97.1	100.0	98.6
98	240	239	1	98.2	97.7	100.0	98.8
99	280	280	0	99.0	98.6	100.0	99.3
99	480	479	1	99.1	98.8	100.0	99.4

^a Table excerpted from Technical Report TR308, *Sampling plans to verify the proportion of an event exceeds or falls below a specified value*, LaBudde, R. (June 4, 2010) (not published). The table was produced as part of an informative report for the Working Group for Validation of Identity Methods for Botanical Raw Materials commissioned by the AOAC INTERNATIONAL Presidential Task Force on Dietary Supplements. The project was funded by the Office of Dietary Supplements, National Institutes of Health.

^b Copyright 2010 by Least Cost Formulations, Ltd. All rights reserved.

^c Based on modified Wilson score 1-sided confidence interval.

^d AOQL = Average outgoing quality level.




Figure A1. Relationship between precision versus bias (trueness). Trueness is reported as bias. Bias is defined as the difference between the test results and an accepted reference value.




Figure A2. Relationship between LOD and LOQ. LOD is defined as the lowest quantity of a substance that can be distinguished from the absence of that substance (a blank value) within a stated confidence limit. LOQ is the level above which quantitative results may be obtained with a stated degree of confidence.




Figure A3. Horwitz Curve, illustrating the exponential increase in the coefficient of variation as the concentration of the analyte decreases [J. AOAC Int. 89, 1095(2006)].

ANNEX B Classification of Methods

The following guidance may be used to determine which performance parameters in Table A1 apply to different classifications of methods. AOAC INTERNATIONAL does not recognize the term “semiquantitative” as a method classification. Methods that have been self-identified as semiquantitative will be classified into one of the following five types:

Type I: Quantitative Methods

Characteristics: Generates a continuous number as a result.

Recommendation: Use performance requirements specified for quantitative method (main or trace component). Use recovery range and maximum precision variation in Tables A4 and A5.

In some cases and for some purposes, methods with less accuracy and precision than recommended in Tables A4 and A5 may be acceptable. Method developers should consult with the appropriate method committee to determine if the recommendations in Tables A4 and A5 do or do not apply to their method.

Type II: Methods that Report Ranges

Characteristics: Generates a “range” indicator such as 0, low, moderate, and high.

Recommendation: Use performance requirements specified for qualitative methods (main component). Specify a range of POD for each range “range” indicator.

Type III: Methods with Cutoff Values

Characteristics: Method may generate a continuous number as an interim result (such as a CT value for a PCR method), which is not reported but converted to a qualitative result (presence/ absence) with the use of a cutoff value.

Recommendation: Use performance requirements specified for qualitative methods.

Type IV: Qualitative Methods

Characteristics: Method of analysis whose response is either the presence or absence of the analyte detected either directly or indirectly in a specified test portion.

Recommendation: Use performance requirements specified for qualitative methods.

Type V: Identification Methods

Characteristics: Method of analysis whose purpose is to determine the identity of an analyte.

Recommendation: Use performance requirements specified for identification methods.

ANNEX C Understanding the POD Model

Excerpted from AOAC INTERNATIONAL Methods Committee Guidelines for Validation of Biological Threat Agent Methods and/or Procedures, J. AOAC Int. 94, 1359(2011) and Official Methods of Analysis of AOAC INTERNATIONAL (2012) 19th Ed., Appendix I.

The Probability of Detection (POD) model is a way of characterizing the performance of a qualitative (binary) method. A binary qualitative method is one that gives a result as one of two possible outcomes, either positive or negative, presence/absence, or +/-.

The single parameter of interest is the POD, which is defined as the probability at a given concentration of obtaining a positive response by the detection method. POD is assumed to be dependent on concentration, and generally, the probability of a positive response will increase as concentration increases.

For example, at very low concentration, the expectation is that the method will not be sensitive to the analyte, and at very high concentration, a high probability of obtaining a positive response is desired. The goal of method validation is to characterize how method response transitions from low concentration/low response to high concentration/high response.




Figure C1. Theoretical POD curve for a qualitative detection method.

POD is always considered to be dependent upon analyte concentration. The POD curve is a graphical representation of method performance, where the probability is plotted as a function of concentration (*see, for example, Figure C1*).

The POD model is designed to allow an objective description of method response without consideration to an a priori expectation of the probabilities at given concentrations. The model is general enough to allow comparisons to any theoretical probability function.

The POD model is also designed to allow for an independent description of method response without consideration to the response of a reference method. The model is general enough to allow for comparisons between reference and candidate method responses, if desired.

Older validation models have used the terms “sensitivity,” “specificity,” “false positive,” and “false negative” to describe method performance. The POD model incorporates all of the performance concepts of these systems into a single parameter, POD.

For example, false positive has been defined by some models as the probability of a positive response, given the sample is truly negative (concentration = 0). The equivalent point on the POD curve for this performance characteristic is the value of the curve at Conc = 0.

Similarly, false negative has sometimes been defined as the probability of a negative response when the sample is truly positive (concentration >0). In the POD curve, this would always be specific to a given sample concentration, but would be represented as the distance from the POD curve to the POD = 1 horizontal top axis at all concentrations except C = 0.

The POD model incorporates all these method characteristics into a single parameter, which is always assumed to vary by concentration. In other models, the terms “false positive,” “false negative,” “sensitivity,” and “specificity” have been defined in a variety of ways, usually not conditional on concentration. For these reasons, these terms are obsolete under this model (*see Table C1*).

The terms “sensitivity,” “specificity,” “false positive,” and “false negative” are obsolete under the POD model (*see Figure C2*).

Table C1. Terminology

Traditional terminology	Concept	POD equivalent	Comment
False positive	Probability of the method giving a (+) response when the sample is truly without analyte	POD(0) POD at conc = 0	POD curve value at conc = 0; “Y-intercept” of the POD curve
Specificity	Probability of the method giving a (-) response when the sample is truly without analyte	1-POD(0)	Distance along the POD axis from POD = 1 to the POD curve value
False negative (at a given concentration)	Probability of a (-) response at a given concentration	1-POD(c)	Distance from the POD curve to the POD = 1 “top axis” in the vertical direction
Sensitivity (at a given concentration)	Probability of a (+) response at a given concentration	POD(c)	Value of the POD curve at any given concentration
True negative	A sample that contains no analyte	C = 0	Point on concentration axis where c = 0
True positive	A sample that contains analyte at some positive concentration	C > 0	Range of concentration where c > 0




Figure C2. Comparison of POD model terminology to other obsolete terms.

ANNEX D
Definitions and Calculations
of HorRat Values from Intralaboratory Data

Excerpted from *Definitions and Calculations of HorRat Values from Intralaboratory Data*, AOAC INTERNATIONAL, *HorRat for SLV.doc*, 2004-01-18.

1. Definitions

1.1 Replicate Data

Data developed under common conditions in the same laboratory: simultaneous performance, or, if necessary to obtain sufficient values, same series, same analyst, same day. Such data provides “repeatability statistical parameters.”

1.2 Pooled Data

Replicate data developed in the same laboratory under different conditions but considered sufficiently similar that, for the purpose of statistical analysis, they may be considered together. These may include different runs, different instruments, different analysts, and different days.

1.3 Average

\bar{x} = Sum of the individual values, x_i , divided by the number of individual values, n .

$$\bar{x} = (\sum x_i) / n$$

1.4 Standard Deviation

$$s_i = [\sum(x_i - \bar{x})^2 / n]^{0.5}$$

1.5 Relative Standard Deviation

$$RSD = s_i \times 100 / \bar{x}$$

1.5.1 Repeatability Relative Standard Deviation [RSD(r) or RSD_r]

The relative standard deviation calculated from within-laboratory data.

1.5.2 Reproducibility Relative Standard Deviation [RSD(R) or RSD_R]

The relative standard deviation calculated from among-laboratory data.

Table D1. Predicted relative standard deviations

Concentration (C)	Mass fraction (C)	PRSD _R , %
100%	1.0	2
1%	0.01	4
0.01%	0.0001	8
1 ppm	0.000001	16
10 ppb	0.00000001	32
1 ppb	0.000000001	45

1.6 Mass Fraction

Concentration, C, expressed as a decimal fraction. For calculating and reporting statistical parameters, data may be expressed in any convenient units (e.g., %, ppm, ppb, mg/g, µg/g; µg/kg; µg/L, µg/µL, etc.). For reporting HorRat values, data must be reported as a mass fraction where the units of the numerator and denominator are the same: e.g., for 100% (pure materials), the mass fraction C = 1.00; for 1 µg/g (ppm), C = 0.000001 = (E-6). See Table D1 for other examples.

1.7 Predicted Relative Standard Deviation [PRSD(R) or PRSD_r]

The reproducibility relative standard deviation calculated from the Horwitz formula:

$$\text{PRSD(R)} = 2C^{-0.15}$$

where C is expressed as a mass fraction. See Table D1.

In spreadsheet notation: PRSD(R) = 2 * C ^(-0.15).

1.8 HorRat Value

The ratio of the reproducibility relative standard deviation calculated from the data to the PRSD(R) calculated from the Horwitz formula:

$$\text{HorRat} = \text{RSD(R)}/\text{PRSD(R)}$$

To differentiate the usual HorRat value calculated from reproducibility data from the HorRat value calculated from repeatability data, attach an R for the former and an r for the latter. But note that the denominator always uses the PRSD(R) calculated from reproducibility data because this parameter is more predictable than the parameter calculated from repeatability data:

$$\text{HorRat(R)} = \text{RSD}_R/\text{PRSD(R)}$$

$$\text{HorRat(r)} = \text{RSD}_r/\text{PRSD(R)}$$

Some expected, predicted relative standard deviations are given in Table D1.

2 Acceptable HorRat Values

2.1 For Interlaboratory Studies

HorRat(R): The original data developed from interlaboratory (among-laboratory) studies assigned a HorRat value of 1.0 with limits of acceptability of 0.5 to 2.0. The corresponding within-laboratory relative standard deviations were found to be typically 1/2 to 2/3 the among-laboratory relative standard deviations.

Table D2. Predicted relative standard deviations

Concentration (C)	PRSD _R , %	PRSD _r , %
100%	2	1
1%	4	2
0.01%	8	4
1 ppm	16	8
10 ppb	32	16
1 ppb	45	22

2.1.1 Limitations

HorRat values do not apply to method-defined (empirical) analytes (moisture, ash, fiber, carbohydrates by difference, etc.), physical properties or physical methods (pH, viscosity, drained weight, etc.), and ill-defined analytes (polymers, products of enzyme reactions).

2.2 For Intralaboratory Studies

2.2.1 Repeatability

Within-laboratory acceptable predicted target values for repeatability are given in Table D2 at 1/2 of PRSD(R), which represents the best case.

2.2.2 HorRat(r)

Based on experience and for the purpose of exploring the extrapolation of HorRat values to SLV studies, take as the minimum acceptability 1/2 of the lower limit ($0.5 \times 0.5 \approx 0.3$) and as the maximum acceptability 2/3 of the upper limit ($0.67 \times 2.0 \approx 1.3$).

Calculate HorRat(r) from the SLV data:

$$\text{HorRat(r)} = \text{RSD(r)}/\text{PRSD(R)}$$

Acceptable HorRat(r) values are 0.3–1.3. Values at the extremes must be interpreted with caution. With a series of low values, check for unreported averaging or prior knowledge of the analyte content; with a series of high values, check for method deficiencies such as unrestricted times, temperatures, masses, volumes, and concentrations; unrecognized impurities (detergent residues on glassware, peroxides in ether); incomplete extractions and transfers and uncontrolled parameters in specific instrumental techniques.

2.3 Other Limitations and Extrapolations

The HorRat value is a very rough but useful summary of the precision in analytical chemistry. It overestimates the precision at the extremes, predicting more variability than observed at the high end of the scale (C > ca 0.1; i.e., >10%) and at the low end of the scale (C < E-8; i.e., 10 ng/g; 10 ppb).

ANNEX E

AOAC Method Accuracy Review

Accuracy of Method Based on Reference Material

Reference material (RM) used.—The use of RMs should be seen as integral to the process of method development, validation, and performance evaluation. RMs are not the only component of a quality system, but correct use of RMs is essential to appropriate quality management. RMs with or without assigned quantity values can be used for measurement precision control, whereas only RMs with assigned quantity values can be used for calibration or measurement trueness control. Method development and validation for matrices within the scope of the method is done to characterize attributes such as recovery, selectivity, “trueness” (accuracy, bias), precision (repeatability and reproducibility), uncertainty estimation, ruggedness, LOQ or LOD, and dynamic range. RMs should be chosen that are fit-for-purpose. When certified reference materials (CRMs) are available with matrices that match the method scope, much of the work involved in method development has already been completed, and that work is documented through the certificate. RMs with analyte values in the range of test samples, as well as “blank” matrix RMs, with values below or near detection limits, are needed.

Availability of RM.—Consideration needs to be given to the future availability of the chosen RM. Well-documented methods that cannot be verified in the future due to lack of material may lose credibility or be seen as inferior.

Fit to method scope.—Natural matrix CRMs provide the greatest assurance that the method is capable of producing accurate results for that matrix. When selecting an RM to perform a method validation, analysts should consider the method to material fit. An example of a good fit would be a method for specified organic molecules in infant formula and using an infant formula or powder milk RM. A poor fit would be a method for specified organic molecules in infant formula and using a sediment material.

Stability.—Providing a stable RM can be challenging where analytes are biologically active, easily oxidized, or interactive with other components of the matrix. CRM producers provide assurance of material stability, as well as homogeneity. CRMs are accompanied by a certificate that includes the following key criteria:

- (1) Assigned values with measurement uncertainty and metrological traceability
- (2) Homogeneity
- (3) Stability, with the expiration date for the certificate
- (4) Storage requirements
- (5) Information on intended use
- (6) Identity of matrix

For some RMs, such as botanical RMs, the source and/or authenticity can be a very important piece of information that should be included with the certificate. Even under ideal storage conditions, many analytes have some rate of change. Recertification may be done by the supplier, and a certificate reissued with a different expiration date and with certain analyte data updated or removed.

Definition of CRM.—Refer to the AOAC TDRM document for definitions from ISO Guide 30, Amd. 1 (2008), <http://www.aoc.org/divisions/References.pdf>.

Information on source of RM is available.—It is the responsibility of the material producer to provide reliable authentication of the RM and make a clear statement in the accompanying documentation. This should be an as detailed listing as possible, including handling of ingredients, identification of plant materials as completely as feasible (species, type, subtype, growing region), etc. This is comparable to other required information on an RM for judging its suitability for a specific application purpose (e.g., containing how much of the targeted analyte, stabilized by adding acid—therefore not suited for certain parameters/procedures, etc.).

Separate RM used for calibration and validation.—A single RM cannot be used for both calibration and validation of results in the same measurement procedure.

Blank RM used where appropriate.—Blank matrix RMs are useful for ensuring performance at or near the detection limits. These are particularly useful for routine quality control in methods measuring, for instance, trace levels of allergens, mycotoxins, or drug residues.

Storage requirements were maintained.—Method developers should maintain good documentation showing that the RM producer’s recommended storage conditions were followed.

Cost.—The cost of ongoing method checks should be considered. Daily use of CRMs can be cost prohibitive. Monthly or quarterly analysis of these materials may be an option.

Concentration of analyte fits intended method.—Concentration of the analyte of interest is appropriate for standard method performance requirements (SMPRs).

Uncertainty available.—Every measurement result has an uncertainty associated with it, and the individual contributions toward the combined uncertainty arise from multiple sources. Achieving the target measurement uncertainty set by the customer for his/her problem of interest is often one of the criteria used in selecting a method for a given application. Estimation of measurement uncertainty can be accomplished by different approaches, but the use of RMs greatly facilitates this part of a method validation.

Demonstration of Method Accuracy when No Reference Material Is Available

If an RM is not available, how is accuracy demonstrated?

There are many analytes for which a CRM with a suitable matrix is not available. This leaves the analyst with few options. For some methods, there may be proficiency testing programs that include a matrix of interest for the analyte. Proficiency testing allows an analyst to compare results with results from other laboratories, which may or may not be using similar methods. Spiking is another technique that may be used. When alternative methods are available, results may be compared between the different methods. These alternatives do not provide the same level of assurance that is gained through the use of a CRM.

Spike recovery.—In the absence of an available CRM, one technique that is sometimes used for assessing performance is the spiking of a matrix RM with a known quantity of the analyte. When this method is used, it cannot be assumed that the analyte is bound in the same way as it would be in a natural matrix. Nevertheless, a certified blank RM would be the preferred choice for constructing a spiked material.

When preparing reference solutions, the pure standards must be completely soluble in the solvent. For insoluble materials in a liquid suspension or for powdered forms of dry materials, validation is required to demonstrate that the analyte is homogeneously distributed and that the response of the detection system to the analyte is not affected by the matrix or preparation technique. When a matrix material is selected for spiking, it should be reasonably

The document, *AOAC Method Accuracy Review*, was prepared by the AOAC Technical Division on Reference Materials (TDRM) and approved by the AOAC Official Methods Board in June 2012.

characterized to determine that it is sufficiently representative of the matrix of interest. Spiked samples must be carried through all steps of the method. Many analytes are bound in a natural matrix and whether the spiked analyte will behave the same as the analyte in a natural matrix is unknown.

Other.—Use of a substitute RM involves the replacement of the CRM with an alternative matrix RM matching the matrix of interest as close as possible based on technical knowledge.

ANNEX F Development and Use of In-House Reference Materials

The use of reference materials is a vital part of any analytical quality assurance program. However, you may have questions about their creation and use. The purpose of this document is to help answer many of these questions.

- What is a reference material?
- Why use reference materials?
- What certified reference materials are currently available?
- Why use an in-house reference material?
- How do I create an in-house reference material?
- How do I use the data from an in-house reference material?

What Is a Reference Material?

The International Organization for Standardization (ISO) defines a reference material as a “material or substance one or more of whose property values are sufficiently homogeneous and well established to be used for the calibration of an apparatus, the assessment of a measurement method, or for assigning values to materials” (1). In plain English, natural-matrix reference materials, such as those you might prepare for use in-house, can be used to validate an analytical method or for quality assurance while you’re using your method to analyze your samples. (Natural-matrix materials are not generally used as calibrants because of the increased uncertainty that this would add to an analysis.) The assigned values for the target analytes of an in-house reference material can be used to establish the precision of your analytical method and, if used in conjunction with a CRM, to establish the accuracy of your method.

ISO defines a certified reference material (CRM) as a “reference material, accompanied by a certificate, one or more of whose property values are certified by a procedure which establishes traceability to an accurate realization of the unit in which the property values are expressed, and for which each certified value is accompanied by an uncertainty at a stated level of confidence” (1).

Why Use Reference Materials?

Certified reference materials can be used across the entire scope of an analytical method and can provide traceability of results to the International System of Units (SI). During method development, CRMs can be used to optimize your method. During method validation, they can be used to ensure that your method is capable of producing the “right” answer, and to determine how close your result is to that answer. During routine use, they can be used to determine within-day and between-day repeatability, and so demonstrate that your method is in control and is producing accurate results every time it is used.

Excerpted from *Development and Use of In-House Reference Materials*, Rev. 2, 2009. Copyright 2005 by the AOAC Technical Division on Reference Materials (TDRM).

Natural-matrix reference materials should mimic the real samples that will be analyzed with a method. They should behave just as your samples would during a procedure, so if you obtain accurate and precise values for your reference material, you should obtain accurate and precise values for your samples as well.

What Certified Reference Materials Are Currently Available?

CRMs are available from a number of sources, including (but not limited to):

- American Association of Cereal Chemists (AACC)
- American Oil Chemists Society (AOCS)
- International Atomic Energy Agency (IAEA)
- Institute for Reference Materials and Measurements (IRMM)
- LGC Promochem
- National Institute of Standards and Technology (NIST)
- National Research Council Canada (NRC Canada)
- UK Food Analysis Proficiency Assessment Program (FAPAS)

A number of websites provide general overviews and catalogs of producers’ and distributors’ reference materials:

- <http://www.aocs.org/tech/crm/>
- <http://www.comar.bam.de>
- <http://www.erm-crm.org>
- <http://www.iaea.org/oregrammes/laqcs>
- <http://www.aaccnet.org/checksample>
- <http://www.irmm-ire.be/mrm.html>
- <http://www.lgcpromochem.com>
- <http://www.naweb.iaea.org/nahu/nmrm/>
- <http://www.nist.gov/srm>
- <http://www.fapas.com/index.cfm>
- <http://www.virm.net>


Because new reference materials are produced regularly, it is important to check these websites to determine what is currently available.

Why Use an In-House Reference Material?

There are many benefits to the use of a CRM. CRMs have been prepared to be homogeneous and, if stored under the proper conditions, stable. You are provided with a certified value as well as the statistical data for the concentration of your analyte; this is about as close as you can come to knowing the true value of the concentration of the analyte. The material has been tested by experienced analysts in leading laboratories, so you have the security of knowing that your method is generating values similar to those generated in other competent laboratories. The CRMs from the sources mentioned above are nationally and/or internationally recognized, so when you obtain acceptable results for a CRM using your analytical method, you give credibility to your methodology and traceability to your results.

But there are some drawbacks associated with CRMs. Unfortunately, many analyte/matrix combinations are not currently available. When testing food products for nutrient content, for example, a laboratory can be asked to analyze anything that might be found in a kitchen or grocery store. Reference materials that represent all of the types of foods that need to be tested are not available, and most CRMs are certified for a limited number of analytes. It is important to match the reference material matrix to your sample matrix. (Food examples dominate the discussion below, but the same processes apply to the development of in-house RMs in other areas of analytical chemistry.)

To demonstrate the applicability of an analytical method to a wide variety of food matrices, AOAC INTERNATIONAL’s Task



Force on Methods for Nutrition Labeling developed a triangle partitioned into sectors in which foods are placed based on their protein, fat, and carbohydrate content (2, 3). Since ash does not have a great impact on the performance of an analytical method for organic-material foods, and water can be added or removed, it can be assumed that the behavior of an analytical method is determined to large extent by the relative proportions of these proximates. AOAC INTERNATIONAL anticipated that one or two foods in a given sector would be representative of other foods in that sector and therefore would be useful for method assessment. Similarly, one or two reference materials in a given sector (or near each other in adjacent sectors) should be useful for quality assurance for analyses involving the other foods in the sector. The positions of many of the food-matrix CRMs from the sources listed above are shown in the triangle and are provided in the list.

These food-matrix reference materials are spread through all sectors of the triangle, thereby making it likely that you can find an appropriate CRM to match to your samples. Ultimately, however, the routine use of a CRM can be cost prohibitive, and is not really the purpose of CRMs. For example, in order to use NIST’s Standard Reference Material (SRM) 2387 Peanut Butter for all mandatory nutrition labeling analyses, you could buy one sales unit (three jars, each containing 170 g material) for \$649 (2009 price). If you charge your customer about \$1000 for analysis of all mandatory nutrients in a test material, the control material would account for more than 60% of your fees. Therefore, many laboratories have found it more cost-effective to create in-house reference materials for routine quality control and characterize them in conjunction with the analysis of a CRM (4). You can prepare larger quantities of a reference material by preparing it in-house, and you have more flexibility in the types of matrices you can use. There are not many limitations on what can be purchased.

How Do I Create an In-House Reference Material?

There are basically three steps to preparing an in-house reference material: selection (including consideration of homogeneity and stability), preparation, and characterization. Additional guidance through these steps can be provided from TDRM as well as in ISO Guides 34 (5) and 35 (6).

References

(1) JCGM 200:2008, *International vocabulary of metrology—Basic and general concepts and associated terms (VIM)*, International Bureau of Weights and Measures (www.bipm.org)

Sector	RM No.	Matrix
	NIST 1563	Coconut oil
1	NIST 3274	Fatty acids in botanical oils
1	NIST 3276	Carrot extract in oil
1	LGC 7104	Sterilized cream
2	NIST 2384	Baking chocolate
3	NIST 2387	Peanut butter
4	NIST 1546	Meat homogenate
4	LGC 7106	Processed cheese
4	LGC 7000	Beef/pork meat
4	LGC 7150	Processed meat
4	LGC 7151	Processed meat
4	LGC 7152	Processed meat
4	SMRD 2000	Fresh meat
4	LGC 7101	Mackerel paste
4	LGC QC1001	Meat paste 1
4	LGC QC1004	Fish paste 1
5	BCR-382	Wheat flour
5	BCR-381	Rye flour
5	LGC 7103	Sweet digestive biscuit
5	LGC 7107	Madeira cake
5	LGC QC1002	Flour 1
6	NIST 1544	Fatty acids
6	NIST 1548a	Typical diet
6	NIST 1849	Infant/adult nutritional formula
6	LGC 7105	Rice pudding
7	LGC 7001	Pork meat
7	NIST 1566b	Oyster tissue
7	NIST 1570a	Spinach leaves
7	NIST 2385	Spinach
8	NIST 1946	Lake trout
8	LGC 7176	Canned pet food
9	NIST 1974a	Mussel tissue
9	NIST 3244	Protein powder

(2) Wolf, W.R., & Andrews, K.W. (1995) *Fresenius’ J. Anal. Chem.* **352**, 73–76

(3) Wolf, W.R. (1993) *Methods of Analysis for Nutrition Labeling*, D.R. Sullivan & D.E. Carpenter (Eds), AOAC INTERNATIONAL, Gaithersburg, MD

(4) European Reference Materials (2005) *Comparison of a Measurement Result with the Certified Value*, Application Note 1

(5) *ISO Guide 34 General Requirements for the Competence of Reference Material Producers* (2009) 2nd, International Organization for Standardization, Geneva, Switzerland

(6) *Guide 35 Certification of Reference Materials—General and Statistical Principles* (2006) International Organization for Standardization, Geneva, Switzerland

For more information about the AOAC Technical Division on Reference Materials, visit <http://aoac.org/divisions/tdrm>.

